


19 
 Europäisches Patentamt
 European Patent Office
 Office européen des brevets

11 Publication number:

0 184 257
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: 85201937.1

22 Date of filing: 22.11.85

61 Int. Cl.⁴: **C 07 D 413/12**

C 07 D 417/12, C 07 D 413/14

C 07 D 417/14, C 07 D 498/04

C 07 D 513/04, A 61 K 31/445

A 61 K 31/42, A 61 K 31/425

30 Priority: 03.12.84 US 677412

43 Date of publication of application:
11.06.86 Bulletin 86/24

84 Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: **JANSSEN PHARMACEUTICA N.V.**
Turnhoutsebaan 30
B-2340 Beerse(BE)


72 Inventor: **Stokbroekx, Raymond Antoine**
Rode Kruisstraat 13
B-2340 Beerse(BE)

72 Inventor: **Luyckx, Marcel Gerebernus Maria**
Eksterstraat 24
B-2440 Geel(BE)

72 Inventor: **Janssens, Frans Eduard**
Tinststraat 79
B-2830 Bonheiden(BE)



54 Benzoxazol- and benzothiazolamine derivatives.

57 Novel benzoxazol- and benzothiazolamine derivatives of the formula



wherein

L represents C₁₋₁₂ alkyl, substituted C₁₋₈ alkyl, aryl C₂₋₆ alkenyl, C₃₋₆ cycloalkyl, substituted C₃₋₆ cycloalkyl, a radical of formula



the pharmaceutically acceptable acid addition salts and possible stereochemically isomeric forms thereof, which compounds are antianoxic and anti-hypoxic agents; pharmaceutical compositions containing such compounds as an active ingredient and methods of preparing said compounds and pharmaceutical compositions.

5

10

BENZOXAZOL- AND BENZOTHAZOLAMINE DERIVATIVES


15 Background of the invention:

In U.S. Patent No. 4,219,559 there are described a number of N-heterocyclyl-4-piperidinamines, wherein said heterocyclyl is imidazole being condensed with benzene or pyridine, which compounds are useful as antihistaminic agents.

20 The compounds of the present invention containing an analogous heterocyclyl radical differ therefrom by the fact that said heterocyclyl is oxazole or thiazole being condensed with benzene or pyridine and by their unexpected anti-anoxic properties.

Description of the preferred embodiments:

25 This invention is concerned with novel benzothiazol- and benzoxazol-amine derivatives which structurally may be represented by the formula



35

the pharmaceutically acceptable acid addition salts and the possible stereochemically isomeric forms thereof, wherein:

$-A^1-A^2-A^3-A^4-$ is a bivalent radical having the formula

- 5 $-CH=CH-CH=CH-$ (a);
 $-N=CH-CH=CH-$ (b);
 $-CH=N-CH=CH-$ (c);
 $-CH=CH-N=CH-$ (d); or
 $-CH=CH-CH=N-$ (e);

10 wherein one or two hydrogen atoms in the radical $-A^1-A^2-A^3-A^4-$ may, each independently from each other, be replaced by halo, C_{1-6} alkyl, hydroxy, C_{1-6} alkyloxy, or trifluoromethyl;

Z is O or S;

n is 0 or the integer 1;



15 R is a member selected from the group consisting of hydrogen, C_{1-6} alkyl, hydroxy and C_{1-6} alkyloxy;

R^1 or R^2 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

20 the dotted line between the nitrogen atom bearing R^1 , the interjacent carbon atom and the nitrogen atom bearing R^2 indicating that a double bond exists between the nitrogen bearing R^1 and said interjacent carbon atom, in which case R^1 is absent, or a double bond exists between the interjacent carbon and the nitrogen bearing R^2 , in which case R^2 is absent; and wherein said interjacent carbon atom is the carbon atom positioned between the nitrogen bearing R^1 and the nitrogen bearing R^2 ;

25 R^3 or R^4 are both hydrogens or R^3 and R^4 combined may form a bivalent radical of formula $-CH_2-CH_2-$;

L is a member selected from the group consisting of C_{1-12} alkyl, substituted C_{1-6} alkyl, aryl C_{2-6} alkenyl, C_{3-6} cycloalkyl, substituted C_{3-6} cycloalkyl, a radical of formula



R^5 being hydrogen or C_{1-12} alkylcarbonyl;

R^6 being hydrogen or C_{1-6} alkyl;


R^7 being aryl, a radical R^8-O or a radical R^8-S ;

B being a bivalent radical of formula $-CH_2-$ or $-O-$; and

5 said R^8 being hydrogen; aryl; 2,3-dihydro-1H-indenyl; benzodioxolyl; (2,3-dihydro-1,4-benzodioxin-2-yl)methyl; (2H-1-benzopyran-2-yl)methyl; or phenyl substituted with C_{2-6} alkenyloxy, C_{1-6} alkylcarbonylamino or C_{1-6} alkylphenylcarbonyl; and

10 wherein said substituted C_{3-6} cycloalkyl is C_{3-6} cycloalkyl being substituted with up to two substituents each independently selected from the group consisting of aryl, aryloxy and cyano;

said substituted C_{1-6} alkyl is C_{1-6} alkyl being substituted with a member selected from the group consisting of C_{3-6} cycloalkyl, pyridinyl, pyridinyloxy, aryl, benzimidazolyl, indolyl, isoxazolyl being optionally
15 substituted with phenyl, 3-oxo-1,2,4-triazolo[4,5-a]pyridin-2(3H)-yl, a radical of formula $ArY-$, a radical of formula R^9O- and a radical of formula



(c);

20

said Ar being aryl;

Y being O, S, NH, $-CONH-$, $-CHCN$, $-CHOR^{10}$ or $-SO_2$;

R^9 being hydrogen, aryl C_{1-6} alkyl, C_{3-6} cycloalkyl, phenyl substituted with phenyl, or phenyl substituted with C_{1-6} alkyloxycarbonyl;

25 R^{10} being hydrogen or C_{1-6} alkyl; and

wherein aryl is phenyl optionally substituted with up to three substituents each independently selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkyloxy, halo, trifluoromethyl, cyano, C_{1-6} alkylcarbonyl, nitro, amino and aminocarbonyl.

30

As used herein the term halo is generic to fluoro, chloro, bromo and iodo; " C_{1-6} alkyl" is meant to include straight and branched saturated hydrocarbon radicals, having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, 1-methylethyl, 1,1-dimethylethyl, propyl, butyl, pentyl, hexyl and the like; C_{1-12} alkyl is meant to include straight
35 and branched saturated hydrocarbon radicals, having from 1 to 12 carbon

atoms; "C₂₋₆ alkenyl" refers to alkenyl radicals having from 2 to 6 carbon atoms, such as, for example, 2-propenyl, 2-butenyl, 3-butenyl, 2-pentenyl and the like; and "C₃₋₆ cycloalkyl" embraces cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

5 Preferred compounds within the invention are those wherein L is aryl C₂₋₆ alkenyl, C₁₋₆ alkyl being substituted with aryloxy, a radical of formula (a) or a radical of formula (b).


Particularly preferred compounds within the invention are those wherein L is as described hereinabove for the preferred compounds and
 10 $-A^1-A^2-A^3-A^4-$ is $-CH=CH-CH=CH-$, n is the integer 1, R² is absent and a double bond exists between the nitrogen bearing R² and the interjacent carbon atom, and R³ and R⁴ are both hydrogen atoms.

More particularly preferred compounds within the invention are those wherein $-A^1-A^2-A^3-A^4-$, n, R² and R³ are as defined hereinabove
 15 for the particularly preferred compounds and wherein a double bond exists between the nitrogen bearing R² and the interjacent carbon atom, and wherein L is C₁₋₆ alkyl substituted with aryloxy, a radical of formula (a) wherein R⁵ and R⁶ are both hydrogen and R⁷ is a radical R⁸-O-, wherein R⁸ is aryl, or L is a radical of formula (b),
 20 wherein B is O, R⁵ and R⁶ are both hydrogen radicals.

Especially preferred compounds within the invention are those particularly preferred compounds wherein R¹ is C₁₋₆ alkyl and aryl is halophenyl.

The most preferred compounds are selected from the group consisting
 25 of 4-[(2-benzothiazolyl)methylamino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol and the pharmaceutically acceptable acid addition salts thereof.


In order to simplify the structural representations of the compounds of formula (I) and of certain precursors and intermediates



represented by the symbol D.

The compounds of formula (I) can generally be prepared by N-alkylating an intermediate of formula (III) with a reagent of formula (II).


5



10 In (II) W represents an appropriate reactive leaving group such as, for example, halo, e.g., chloro, bromo or iodo, or a sulfonyloxy group, e.g. methylsulfonyloxy or 4-(methylphenyl)sulfonyloxy.

The alkylation reaction is conveniently conducted in an inert organic solvent such as, for example, an aromatic hydrocarbon, e.g., benzene, methylbenzene, dimethylbenzene, and the like; a C_{1-6} alkanol, e.g., 15 methanol, ethanol, 1-butanol and the like; a ketone, e.g., 2-propanone, 4-methyl-2-pentanone and the like; an ether, e.g., 1,4-dioxane, 1,1'-oxybisethane, tetrahydrofuran and the like; N,N-dimethylformamide (DMF); N,N-dimethylacetamide (DMA); dimethyl sulfoxide (DMSO); nitrobenzene; 1-methyl-2-pyrrolidinone; and the like. The addition of an 20 appropriate base such as, for example, an alkali metal carbonate or hydrogen carbonate, sodium hydride or an organic base such as, for example, N,N-diethylethanamine or N-(1-methylethyl)-2-propanamine may be utilized to pick up the acid which is liberated during the course of the reaction. In some circumstances the addition of an iodide salt, 25 preferably an alkali metal iodide, is appropriate. Somewhat elevated temperatures may enhance the rate of the reaction.


The compounds of formula (I) can also be prepared by the reductive N-alkylation reaction of (III) with an appropriate carbonyl-compound of 30 formula $\text{L}'=\text{C}=\text{O}$ (IV), said $\text{L}'=\text{C}=\text{O}$ being a compound of formula L-H wherein a $-\text{CH}_2-$ radical is oxidated to a carbonyl radical.



35

Said reductive N-alkylation reaction may conveniently be carried out by catalytically hydrogenating a stirred and heated mixture of the reactants in a suitable reaction-inert organic solvent according to art-known catalytic hydrogenating procedures. The reaction mixture may be stirred and/or heated in order to enhance the reaction rate. Suitable solvents are, for example, water; C₁₋₆ alkanols, e.g. methanol, ethanol, 2-propanol and the like; cyclic ethers, e.g. 1,4-dioxane and the like; halogenated hydrocarbons, e.g. trichloromethane and the like; N,N-dimethylformamide; dimethyl sulfoxide and the like; or a mixture of 2 or more of such solvents. The term "art-known catalytic hydrogenating procedures" means that the reaction is carried out under hydrogen atmosphere and in the presence of an appropriate catalyst such as, for example, palladium-on-charcoal, platinum-on-charcoal and the like. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products it may be advantageous to add an appropriate catalyst-poison to the reaction mixture, e.g., thiophene and the like.





The compounds of formula (I) can also be prepared by oxidatively cyclizing an urea or thiourea derivative of formula (V) in the presence of an appropriate oxidant.



Appropriate oxidants are, for example, halogens, e.g. chlorine and bromine, thionyl chloride, sulfuryl chloride, thionyl bromide, sulfuryl bromide and the like agents. The said cyclization reaction is most conveniently conducted in a reaction-inert solvent such as, for example, an halogenated hydrocarbon, e.g. tetrachloromethane, trichloromethane and the like. Elevated temperatures may be appropriate to enhance the reaction rate.

The compounds of formula (I) wherein L is a radical of formula (a) or (b) wherein R⁵ is hydrogen can also be prepared by reacting an

appropriate oxirane derivative of formula (VI), respectively of formula (VIII) with (III). The thus obtained compounds of formula (I-a-1) respectively of formula (I-a-3) may further be converted into the corresponding compounds of formula (I-a-2), respectively (I-a-4) by an appropriate acylation reaction.





The reagent R^{5-a}W^1 represents an alkanolic acid or a reactive derivative thereof such as, for example, an ester, an acid halide or anhydride and the like.

The reaction of (VI) or (VIII) with (III) may conveniently be conducted in an appropriate solvent such as, for example, an aromatic hydrocarbon, e.g. benzene, methylbenzene and the like, a C_{1-6} alkanol, e.g. methanol and ethanol. Elevated temperatures may be appropriate to enhance the reaction rate.

The compounds of formula (I), wherein R^2 is absent and a double bond exists between the nitrogen bearing R^2 and the previously defined interjacent carbon atom, said compounds being represented by the formula (I-b) can also be prepared by reacting an appropriate reagent of formula

5 (IX) with a benzothiazole or benzoxazole of formula (X).




In (IX) and (X) Q^1 and Q^2 are selected so that during the alkylation


20 reaction a radical of formula $-NR^1$ is formed. For example, where Q^1 is an appropriate leaving group, Q^2 is a radical of formula $-NHR^1$, or where Q^1 is a radical of formula $-NHR^1$, Q^2 is an appropriate leaving group.

The compounds of formula (I-b) wherein Z is O and R^1 is hydrogen,

25 said compounds being represented by the formula (I-b-1), can also be prepared by cyclodesulfurizing an intermediate of formula (XIV), which may conveniently be formed in situ by condensing an isothiocyanate (XII) with an aromatic aminoalcohol of formula (XIII).



35




10 Said cyclodesulfurization reaction may be carried out by the reaction of (XIV) with an appropriate alkyl halide, preferably iodomethane in an appropriate reaction-inert organic solvent, e.g., a C₁₋₆ alkanol such as methanol, ethanol, 2-propanol and the like. Otherwise, the cyclodesulfurization reaction may be carried out by the reaction of (XIV) with an appropriate metal oxide or salt in an appropriate solvent according to art-known procedures. For example, the compounds of formula

15 (I) can easily be prepared by the reaction of (XIV) with an appropriate Hg(II) or Pb(II) oxide or salt, such as, for example HgO, HgCl₂, Hg(OAc)₂, PbO or Pb(OAc)₂. In certain instances it may be appropriate to supplement the reaction mixture with a small amount of sulfur. Even so methanediimines, especially N,N'-methanetetraylbis-

20 [cyclohexanamine] may be used as cyclodesulfurizing agents.


The compounds of formula (I-b) wherein R¹ is hydrogen, said compounds being represented by the formula (I-b-2), may be prepared by

25 cyclizing an intermediate (XVII), which in situ may be formed by reacting an isocyanate or isothiocyanate (XV) with an aromatic amine (XVI).



In (XVI) and (XVII) W^2 represents an appropriate leaving group such as, for example, halo, e.g. chloro and bromo; C_{1-6} alkyloxy and C_{1-6} alkylthio.

5 The compounds of formula (I) wherein L is a C_{1-12} alkyl radical or a substituted C_{1-6} alkyl radical, wherein said C_{1-12} alkyl or C_{1-6} alkyl contains at least two carbon atoms, said compounds being represented by the formula (I-c), may also be prepared by reacting an alkene of formula (XVIII) with an intermediate of formula (III) by stirring and, if desired, heating the reactants together, preferably in
10 a suitable solvent such as, for example, an alcohol, e.g. methanol, ethanol, butanol and the like.



15 In (XVIII) R^{11} represents C_{1-6} alkyl, C_{3-6} cycloalkyl, pyridinyl, pyridinyloxy, aryl, benzimidazolyl, indolyl, isoxazolyl being substituted with phenyl, 3-oxo-1,2,4-triazolo[4,5-a]pyridin-2(3H)-yl, a radical of formula ArY , a radical of formula R^9O and a radical of formula (g); and in (I-c) r is an integer of from 2 to 6 included.

20 Additionally, the compounds of formula (I), wherein L is a radical of formula $Ar-CO-NH-CH_2-CH_2-$, said compounds being represented by the formula (I-d), can also be prepared by reacting an aziridine of formula (XIX) with (III), preferably in the presence of a suitable solvent such as, for example, an aromatic hydrocarbon, e.g. benzene,
25 methylbenzene and the like.



30 The compounds of formula (I) can also be converted into each other following art-known procedures. The compounds of formula (I) wherein R^1 is hydrogen may be converted into compounds of formula (I) wherein R^1 is other than hydrogen by reacting the former compounds with an appropriate N-alkylating reagent.

35 The compounds of formula (I) wherein L is a C_{1-6} alkyl radical

substituted with hydroxy may further be Q-alkylated with an appropriate reagent following art-known Q-alkylating procedures.

In all of the foregoing and in the following preparations, the reaction products may be isolated from the reaction mixture and, if
5 necessary, further purified according to methodologies generally known in the art.

The compounds of formula (I) have basic properties and, consequently, they may be converted to their therapeutically active non-toxic acid addition salt forms by treatment with appropriate acids,
10 such as, for example, inorganic acids, such as hydrohalic acid, e.g. hydrochloric, hydrobromic and the like, and sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-hydroxy-propanoic, 2-oxopropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic,
15 (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, benzoic, 2-hydroxybenzoic, 3-phenyl-2-propenoic, α-hydroxybenzeneacetic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form
20 can be converted by treatment with alkali into the free base form.

A number of intermediates and starting materials in the foregoing preparations are known compounds which may be prepared according to art-known methodologies of preparing said or similar compounds. A number
25 of such preparation methods will be described hereinafter in more detail.

The intermediates of formula (II) wherein L is C₁₋₆ alkyl being substituted with pyridinyloxy, aryloxy, arylthio, arylamino or a radical R^{9-a}O-, said R^{9-a} having the previously defined meaning of R⁹,
provided that R^{9-a} is other than hydrogen, can be prepared by
30 alkylating an appropriate alcohol, thioalcohol or amine with a reagent W-C₁₋₆ alkyl-W, or alternatively with a reagent W-Alk-OH, and subsequently converting the thus formed alcohol with a reagent capable of converting an alcohol function into a reactive leaving group, e.g. thionyl chloride, methylsulfonyl chloride and the like.

35 The intermediates of formula (III) can generally be prepared by


5




15


20





The intermediates of formula (III) wherein R^2 is absent and a double bond exists between the nitrogen bearing said R^2 and the previously defined interjacent carbon atom, said intermediates being represented by the formula (III-b) can also be prepared by cyclizing an intermediate (XXVI) which in situ may be formed by reacting an isocyanate or isothiocyanate (XXV) with (XVI), and subsequently eliminating the protective group P as described hereinabove.






10

The previously described intermediates can also be converted into each other following art-known functional group transformation procedures.


For example, the intermediates of formula (V) wherein R^1 is hydrogen and the intermediates of formula (V) wherein R^2 is hydrogen, said intermediates being represented by the formula (V-a), respectively (V-b) can be prepared by reacting an isocyanate or isothiocyanate (XXVIII) with an aromatic amine (XXIX), respectively by reacting an amine (XXX) with an aromatic isocyanate or isothiocyanate (XXXI).

20





30

35



15 The intermediates of formula (VI) and of formula (VII) may be prepared by epoxidizing an alkene reagent of formula (XXXII) respectively of formula (XXXIII) with an appropriate epoxidizing reagent, e.g. with an aromatic peroxoacid, in a suitable solvent.



The intermediates of formula (XIX) may be prepared following the procedures described hereinabove for the preparation of (V-a) and (V-b).

During one of the reactions the intermediates wherein R^1 is hydrogen may be converted into the corresponding intermediates wherein R^1 is other than hydrogen following art-known N-alkylating procedures.

30

From formula (I) it is evident that the compounds of this invention may have several asymmetric carbon atoms in their structure. Each of these chiral centers may be present in a R- and a S-configuration, this R- and S-notation being in correspondence with the rules described by

35

R.S. Cahn, C. Ingold and V. Prelog in Angew. Chem., Int. Ed. Engl., 5, 385, 511 (1966).

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures.

5 Diastereoisomers may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g., counter current distribution, and enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids.

10 Pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically.

It is evident that the cis and trans diastereomeric racemates may be further resolved into their optical isomers, cis(+), cis(-), trans(+)
15 and trans(-) by the application of methodologies known to those skilled in the art.

Stereochemically isomeric forms of the compounds of formula (I) are naturally intended to be embraced within the scope of the invention.

20 The compounds of formula (I), their pharmaceutically acceptable acid-addition salts and stereochemically isomeric forms possess useful pharmacological properties. For example, they show useful anti-hypoxic and anti-anoxic properties, which activity is clearly evidenced by the experimental data obtained in e.g. the "KCN Test in Rats".

25 In view of their anti-anoxic or anti-hypoxic activity, the compounds of formula (I), their acid addition salts and/or their possible stereoisomers are very useful in the treatment of warm-blooded animals suffering from anoxia or hypoxia, for example, in such situations where the oxygen supply is reduced or blocked, such as shock, cardiac arrest,
30 severe blood losses and the like situations.

In view of their anti-anoxic and anti-hypoxic activity the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, in base
35 or acid-addition salt form, as the active ingredient is combined in

intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions: or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed.

Acid addition salts of (I) due to their increased water solubility over the corresponding base form, are obviously more suitable in the preparation of aqueous compositions.

25

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers,

35

injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

5 The present invention is also related with a method of treating hypoxia or anoxia in warm-blooded animals suffering from said hypoxia or anoxia by administering an effective anti-anoxic or hypoxic amount of a compound of formula (I) or a pharmaceutically acceptable acid addition salt thereof.

10 Although the amounts to be administered to subjects may vary within rather large limits, daily doses varying of from 0.1 mg/kg body weight to 40 mg/kg body weight are preferred.

15 The following examples are intended to illustrate and not to limit the scope of the present invention. Unless otherwise stated all parts therein are by weight.

20

25

30

35

EXPERIMENTAL PARTA. Preparation of IntermediatesExample 1

5 A mixture of 100 parts of ethyl 4-oxo-1-piperidinecarboxylate, 42 parts of 2-propanamine, 2 parts of a solution of thiophene in methanol 4% and 400 parts of methanol was hydrogenated at normal pressure and at room temperature with 5 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off and the filtrate was
10 evaporated, yielding 118 parts (94%) of ethyl 4-[(1-methylethyl)-amino]-1-piperidinecarboxylate (1).

In a similar manner there were also prepared:

- ethyl 3-(methylamino)-8-azabicyclo[3.2.1]octane-8-carboxylate (2);
ethyl 3-(methylamino)-1-piperidinecarboxylate (3);
15 cis-ethyl 3-(phenylmethoxy)-4-[(phenylmethyl)amino]-1-piperidinecarboxylate (4);
ethyl 4-(butylamino)-1-piperidinecarboxylate (5);
cis-ethyl 3-methoxy-4-(methylamino)-1-piperidinecarboxylate monohydrochloride; mp. 169.1°C (6);
20 cis-ethyl 4-(methylamino)-3-(phenylmethoxy)-1-piperidinecarboxylate monohydrochloride; mp. 181.5°C (7);
ethyl (cis+trans)-3-methyl-4-(methylamino)-1-piperidinecarboxylate (8); and
ethyl 3-(methylamino)-1-pyrrolidinecarboxylate (9).

25 Example 2

A mixture of 150 parts of cis-ethyl 4-(methylamino)-3-(phenylmethoxy)-1-piperidinecarboxylate and 560 parts of methanol was hydrogenated at normal pressure and at 50°C with 10 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of
30 hydrogen was taken up, the catalyst was filtered off and the filtrate was evaporated, yielding 101 parts (95%) of ethyl cis-3-hydroxy-4-(methylamino)-1-piperidinecarboxylate (10).

In a similar manner there was also prepared:

- cis-ethyl 4-amino-3-hydroxy-1-piperidinecarboxylate; bp. 175-185°C at
35 0.4 mm pressure (11).

Example 3

A mixture of 8.6 parts of sodium hydroxide and 120 parts of water was stirred at a temperature below 10°C and there were added successively 17 parts of carbon disulfide and 35 parts of ethyl 3-amino-1-pyrrolidinecarboxylate. Stirring was continued for 3 hours. Then there were added dropwise 23.5 parts of ethyl carbonylchloridate. Upon completion, stirring was continued for 2 hours at 60°C. The reaction mixture was extracted with methylbenzene. The extract was dried, filtered and evaporated, yielding 55 parts (100%) of ethyl 3-isothiocyanato-1-pyrrolidinecarboxylate (12).

In a similar manner there was also prepared:
ethyl 4-isothiocyanato-1-piperidinecarboxylate (13).

Example 4


A mixture of 107 parts of ethyl 4-isothiocyanato-1-piperidinecarboxylate, 53.5 parts of *N*-methylbenzenamine, 1 part of *N,N*-dimethyl-4-pyridinamine and 450 parts of methylbenzene was stirred and refluxed overnight. After cooling, the reaction mixture was evaporated. The residue was crystallized from 2,2'-oxybispropane. The product was filtered off and dried, yielding 68 parts (42%) of ethyl 4-[[[(methylphenylamino)thioxomethyl]amino]-1-piperidinecarboxylate; mp. 95.2°C (14).

In a similar manner there were also prepared:
ethyl 4-[(phenylamino)thioxomethylamino]-1-piperidinecarboxylate (15); and
ethyl 4-[[[(2-hydroxyphenyl)aminothioxomethyl]amino]-1-piperidinecarboxylate; mp. 130°C (16).

Example 5

To a stirred solution of 91 parts of ethyl 4-(methylamino)-1-piperidinecarboxylate in 420 parts of 2,2'-oxybispropane were added dropwise 75 parts of 1-fluoro-3-isothiocyanatobenzene. Upon completion, stirring was continued for 1 hour. The precipitated product was filtered off and dried, yielding 150 parts (90%) of ethyl 4-[[[(3-fluorophenyl)aminothioxomethyl]methylamino]-1-piperidinecarboxylate; mp. 175.2°C (17).

In a similar manner there were also prepared:



5

No.	R ¹	R'	R	isomerism	mp.(°C)
18	H	H	OH	cis	191.1
19	CH ₃	H	H	-	166.0
20	C ₂ H ₅	H	H	-	oil
21	n.C ₄ H ₉	H	H	-	-
22	CH ₃	4-CH ₃	H	-	-
23	C ₆ H ₅ CH ₂	H	H	-	-
24	CH ₃	4-F	H	-	178.6
25	CH ₃	H	OCH ₃	cis	166.0
26	CH ₃	3-OH	H	-	-
27	CH ₃	H	CH ₃	cis+trans	-
28	1.C ₃ H ₇	H	H	-	-
29	CH ₃	3-CH ₃ O	H	-	-
30	CH ₃	3-Cl	H	-	-
31	CH ₃	4-OH	H	-	200.0
32	CH ₃	4-Cl	H	-	193.1
33	CH ₃	4-CH ₃ O	H	-	-
34	CH ₃	H	OH	cis	-

25

In a similar manner there were also prepared:

ethyl 3-[methyl[(phenylamino)thioxomethyl]amino]-8-azabicyclo-[3.2.1]octane-8-carboxylate; mp. 143.9°C (35);

30

ethyl 3-[methyl[(phenylamino)thioxomethyl]amino]-1-piperidine-carboxylate (36); and

ethyl 3-[methyl[(phenylamino)thioxomethyl]amino]-1-pyrrolidine-carboxylate; mp. 145.4°C (37).

35

Example 6

A mixture of 55 parts of ethyl 3-isothiocyanato-1-pyrrolidine-carboxylate, 23.6 parts of 2-aminophenol and 320 parts of acetonitrile was stirred and refluxed for 4 hours. Then there were added
5 93 parts of mercury(II) oxide and 0.5 parts of sulfur while stirring vigorously. The reaction mixture was stirred and refluxed overnight. After cooling, the mixture was filtered over diatomaceous earth and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane
10 and methanol (99:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 38 parts (64%) of ethyl 3-[(2-benzoxazolyl)amino]-1-pyrrolidinecarboxylate as a residue (38).

In a similar manner there were also prepared:

15 ethyl 4-(2-benzoxazolylamino)-1-piperidinecarboxylate monohydrobromide (39).

ethyl 4-[(oxazolo[4,5-b]pyridin-2-yl)amino]-1-piperidinecarboxylate monohydrobromide; mp. 203.0°C (40).


In a similar manner there is also prepared:

20 ethyl 4-[(oxazolo[4,5-d]pyridin-2-yl)amino]-1-piperidinecarboxylate (41).

Example 7

To a stirred mixture of 196 parts of ethyl 4-[ethyl[(phenyl-amino)thioxomethyl]amino]-1-piperidinecarboxylate and 1600 parts of
25 tetrachloromethane were added dropwise 89 parts of bromine at room temperature. Upon completion, stirring was continued first for 30 minutes at room temperature and further for 4 hours at reflux. The reaction mixture was cooled and the tetrachloromethane was decanted, yielding 194 parts (100%) of ethyl 4-[(2-benzothiazolyl)ethyl-
30 amino]-1-piperidinecarboxylate as an oily residue (42).

In a similar manner there were also prepared:



5

No.	R ¹	R'	R	R''	isomerism	salt/base	mp. (°C)
43	H	H	H	H	-	HBr	-
44	CH ₃	H	H	H	-	HBr	-
45	n.C ₄ H ₉	H	H	H	-	HBr	oil
46	CH ₃	CH ₃	H	H	-	base	-
47	C ₆ H ₅ CH ₂	H	H	H	-	base	-
48	CH ₃	H	OCH ₃	H	cis	base	-
49	CH ₃	H	H	F	-	base	oil
50	CH ₃	H	H	OH	-	HBr	-
51	CH ₃	H	CH ₃	H	cis+trans	HBr	-
52	1.C ₃ H ₇	H	H	H	-	HBr	-
53	CH ₃	H	H	OCH ₃	-	base	-
54	CH ₃	H	H	Cl	-	HBr	-
55	CH ₃	OH	H	H	-	HBr	-
56	CH ₃	Cl	H	H	-	HBr	-
57	CH ₃	H	OH	H	cis	base	-
58	H	H	OH	H	cis	base	-

25

In a similar manner there were also prepared:

ethyl 3-[(2-benzothiazolyl)methylamino]-8-azabicyclo[3.2.1]octane-8-carboxylate (59);

ethyl 3-[(2-benzothiazolyl)methylamino]-1-piperidinecarboxylate (60);

ethyl 3-[(2-benzothiazolyl)methylamino]-1-pyrrolidinecarboxylate

30

monohydrobromide (61); and

ethyl 4-[(3-methyl-2(3H)-benzothiazolylidene)amino]-1-piperidine-carboxylate (62).

35

Example 8

To a stirred mixture of 103.1 parts of ethyl 4-[[[(3-chloro-phenyl)amino]thioxomethyl]methylamino]-1-piperidinecarboxylate and 960 parts of tetrachloromethane were added dropwise 45.4 parts of
5 bromine at about 20°C. Upon completion, stirring was continued first for 45 minutes at room temperature and further for 35 hours at reflux. The reaction mixture was cooled and the tetrachloromethane was decanted, yielding 126 parts (100%) of ethyl 4-[(7-chloro-2-benzothiazolyl)methylamino]-1-piperidinecarboxylate monohydrobromide
10 as a residue (63):

Example 9

To a stirred mixture of 124 parts of ethyl 4-[[[(4-methoxy-phenyl)amino]thioxomethyl]methylamino]-1-piperidinecarboxylate and 1500 parts of tetrachloromethane were added dropwise slowly 112.6
15 parts of bromine at room temperature. Upon completion, stirring was continued first for 30 minutes at room temperature and then for 3 hours at reflux temperature. After cooling, tetrachloromethane was decanted, yielding 110 parts (73%) of ethyl 4-[(5-bromo-6-methoxy-2-benzothiazolyl)methylamino]-1-piperidinecarboxylate as a residue
20 (64).

Example 10

A mixture of 51 parts of ethyl 4-isothiocyanato-1-piperidinecarboxylate, 30 parts of 2-chloro-3-pyridinamine and 240 parts of ethanol was stirred and refluxed for 24 hours. The reaction mixture
25 was cooled to room temperature. Upon the addition of 70 parts of 2,2'-oxybispropane, the product was allowed to crystallize. It was filtered off and dried, yielding 40 parts (50%) of ethyl 4-(thiazolo[5,4-b]pyridin-2-ylamino)-1-piperidinecarboxylate monohydrochloride (65).

Example 11

A mixture of 8.5 parts of ethyl 4-amino-1-piperidinecarboxylate and 2.6 parts of 2-(methylsulfonyl)thiazolo[4,5-c]pyridine was molten together for 1 hour at 120°C. After cooling, the mixture was taken up in trichloromethane. The whole was washed with water and
35 sodium hydroxide, dried, filtered and evaporated. The residue was


purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (99:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was stirred in warm 2,2'-oxybispropane. The product was filtered off and dried, yielding 2.8 parts (76%) of ethyl 4-[(thiazolo[4,5-c]-pyridin-2-yl)amino]-1-piperidinecarboxylate; mp. 213°C (66).

Example 12

A mixture of 20 parts of ethyl 4-[(2-benzothiazolyl)methylamino]-1-piperidinecarboxylate monohydrobromide and 300 parts of hydrobromic acid solution 48% in water was stirred and refluxed for 2 hours. The reaction mixture was evaporated and the residue was recrystallized from methanol, yielding 17 parts (85%) of N-methyl-N-(4-piperidiny)-2-benzothiazolamine dihydrobromide; mp. + 260°C (67).

A solution of 427 parts of N-methyl-N-(4-piperidiny)-2-benzothiazolamine dihydrobromide monohydrate in 1500 parts of water was treated with a sodium hydroxide solution 50%. The product was extracted twice with trichloromethane. The combined organic layers were washed with water, dried, filtered and evaporated. The residue was crystallized from 490 parts of 2,2'-oxybispropane. The product was filtered off and dried, yielding 248 parts (100%) of N-methyl-N-(4-piperidiny)-2-benzothiazolamine; mp. 86.9°C (68).

In a similar manner there were also prepared:



25

No.	R	R ¹	A ¹ =A ² -A ³ =A ⁴	Z	isomerism	salt/base	mp.(°C)
69	H	H	CH=CH-CH=CH	S	-	2HBr	260
70	OH	H	CH=CH-CH=CH	S	cis	2HBr.H ₂ O	268-300
71	H	H	N=CH-CH=CH	S	-	3HBr	286.9
72	H	CH ₃	N=CH-CH=CH	S	-	base	144
73	H	H	CH=CH-CH=CH	O	-	2HBr	263
74	H	CH ₃	CH=CH-CH=CH	O	-	2HBr.1/2H ₂ O	261.2
75	H	CH ₃	CH=CH-CH=CH	O	-	base	130.3

30

35

5	76	H	C ₂ H ₅	CH=CH-CH=CH	S	-	*	191.7
	77	H	n.C ₄ H ₉	CH=CH-CH=CH	S	-	**	160.0
	78	H	CH ₃	CH=C-CH=CH CH ₃	S	-	base	74.3
	79	H	C ₆ H ₅ CH ₂	CH=CH-CH=CH	S	-	*	206.4
	80	OCH ₃	CH ₃	CH=CH-CH=CH	S	cis	(COOH) ₂	206.4
10	81	H	CH ₃	CH=CH-CF=CH	S	-	2HCl.H ₂ O	266.4
	82	H	CH ₃	CH=CF-CH=CH	S	-	base	-
	83	H	CH ₃	CH-CH-C-CH OH	S	-	2HBr	-
	84	H	1.C ₃ H ₇	CH=CH-CH=CH	S	-	*	181
	85	H	CH ₃	CH-CH-C-CH OCH ₃	S	-	base	-
15	86	H	CH ₃	CH=CH-C-CH Cl	S	-	*	158
	87	H	CH ₃	C=CH-CH=CH Cl	S	-	base	-
	88	H	CH ₃	CH=C-CH=CH OH	S	-	2HBr	-
	89	H	CH ₃	CH-C-CH=CH Cl	S	-	2HCl.H ₂ O	237.8
	90	H	CH ₃	CH=C-CBr=CH OCH ₃	S	-	base	137.0
25	91	OH	CH ₃	CH=CH-CH=CH	S	cis	base	-

* : (Z)-2-butenedioate (1:1)

** : (Z)-2-butenedioate (2:3)

In a similar manner there were also prepared:

30 N-(1-methylthiazolo[5,4-b]pyridin-2(1H)-ylidene)-4-piperidinamine (92);

N-(2-benzothiazolyl)-N-methyl-8-azabicyclo[3.2.1]octan-3-amine (E)-2-butenedioate(1:1); mp. 236.1°C (93);

N-methyl-N-(3-piperidiny)-2-benzothiazolamine (94);

N-methyl-N-(3-pyrrolidinyl)-2-benzothiazolamine ethanedioate(1:1);
mp. 157.0°C (95);

N-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine; mp. 198.5°C
(96);

5 N-methyl-N-(3-pyrrolidinyl)-2-benzoxazolamine (97);

N-(4-piperidinyl)thiazolo[4,5-c]pyridin-2-amine (98); and

N-(4-piperidinyl)oxazolo[4,5-b]pyridin-2-amine (99).

In a similar manner there is also prepared:

N-(4-piperidinyl)oxazolo[4,5-d]pyridin-2-amine (100).

10 Example 13

A mixture of 100 parts of ethyl (cis+trans)-4-[(2-benzothiazolyl)-methylamino]-3-methyl-1-piperidinecarboxylate monohydrobromide and 600 parts of a hydrobromic acid solution 48% in water was stirred and refluxed for 3 hours. After cooling, the reaction mixture was
15 evaporated. The residue was crystallized from 2-propanol : two fractions were obtained. The first fraction was taken up in water. The solution was treated with sodium hydroxide. The product was extracted with trichloromethane. The organic layer was washed with water, dried, filtered and evaporated, yielding 30 parts (48%) of
20 (cis+trans)-N-methyl-N-(3-methyl-4-piperidinyl)-2-benzothiazolamine (101) as an oily residue. The second fraction was taken up in water. The solution was treated with sodium hydroxide. The product was extracted with trichloromethane. The organic layer was washed with water, dried, filtered and evaporated. The residue was converted
25 into the hydrobromide salt in methanol. The salt was filtered off and dried, yielding 16.6 parts (16%) of cis-N-methyl-N-(3-methyl-4-piperidinyl)-2-benzothiazolamine dihydrobromide; mp. 266.7°C (102).

Example 14

To a stirred mixture of 38 parts of ethyl 3-[(2-benzoxazolyl)-amino]-1-pyrrolidinecarboxylate, 150 parts of dimethyl sulfoxide and
30 135 parts of benzene were added portionwise 7.25 parts of a sodium hydride dispersion 50% at a temperature below 10°C. After stirring for a while, 21.6 parts of iodomethane were added dropwise at the same temperature. Upon completion, stirring was continued overnight
35 at 50°C. 1000 Parts of water were added and the product was

extracted with methylbenzene. The extract was washed with water, dried, filtered and evaporated. The residue was converted into the hydrochloride salt in 2-propanol. The salt was filtered off and dried, yielding 40 parts (80%) of ethyl 3-[(2-benzoxazolyl)methylamino]-1-pyrrolidinecarboxylate dihydrochloride (103).

In a similar manner there were also prepared:

ethyl 4-[(2-benzoxazolyl)methylamino]-1-piperidinecarboxylate monohydrobromide; mp. 152.6°C (104);

ethyl 4-(1-methylthiazolo[5,4-b]pyridin-2(1H)-ylidenamino)-1-piperidinecarboxylate (105); and
ethyl 4-[methyl(thiazolo[5,4-b]pyridin-2-yl)amino]-1-piperidinecarboxylate (106).

Example 15

To a stirred mixture of 32.1 parts of 3-fluorophenol, 122 parts of 1,6-dibromohexane and 200 parts of water was dropwise added a solution of 20 parts of sodium hydroxide in 100 parts of water at reflux temperature. Upon completion, stirring was continued for 24 hours at reflux. After cooling, the product was extracted three times with trichloromethane. The combined extracts were dried, filtered and evaporated. The residue was distilled, yielding 40 parts (50%) of 1-[(6-bromohexyl)oxy]-3-fluorobenzene; bp. 45-90°C at 0.05 mm pressure (107).

In a similar manner there were also prepared:



25

No.	Ar	Y	n	W	physical data
108	4-C ₆ H ₅ -C ₆ H ₄	O	3	Cl	mp. 61.5-63°C
109	2-CH ₃ CO, 4F-C ₆ H ₃	O	3	Cl	bp. 130-147°C (0.2 mm Hg)
110	4-CH ₃ OCO-C ₆ H ₄	O	4	Cl	bp. 157°C (0.4 mm Hg)
111	2F-C ₆ H ₄	O	5	Br	bp. 110-112°C (0.6 mm Hg)
112	4-C ₂ H ₅ -C ₆ H ₄	O	5	Br	bp. 128-134°C (0.2 mm Hg)
113	3-CH ₃ O-C ₆ H ₄	O	5	Br	bp. 150-154°C (0.4 mm Hg)
114	1-naphthalenyl	O	5	Br	bp. 165-170°C (0.6 mm Hg)
115	3-Cl-C ₆ H ₄	O	5	Br	bp. 135-137°C (0.7 mm Hg)

	116	$3\text{-CH}_3\text{CO-C}_6\text{H}_4$	O	3	Cl	bp. 120°C (0.5 mm Hg)
	117	$3,4,5(\text{Cl})_3\text{-C}_6\text{H}_2$	O	3	Br	-
	118	$2\text{-NH}_2\text{CO}_2\text{F-C}_6\text{H}_2$	O	3	Cl	mp. 108°C
5	119	$4\text{-F-C}_6\text{H}_4$	S	3	Br	bp. 78-80°C (0.2 mm Hg)
	120	$2,6\text{-(Br)}_2,4\text{CH}_3\text{-C}_6\text{H}_2$	O	3	Br	bp. 118-123 (0.3 mm Hg)
	121	1-Br-2-naphthalenyl	O	3	Br	-
	122	$3\text{-F-C}_6\text{H}_4$	O	5	Br	bp. 120°C (0.1-0.30 mm Hg)
	123	$3\text{-F-C}_6\text{H}_4$	O	4	Br	-
10	124	$4\text{-CH}_3\text{CO-C}_6\text{H}_4$	O	5	Br	bp. 165-170°C (0.02 mm Hg)

And following the same procedures and using the appropriate starting materials there were also prepared:

[(6-bromo-2-naphthalenyloxy)methyl]oxirane (125);

15 (4-methylphenyl)[4-(oxiranylmethoxy)phenyl]methanone (126);

[(2,6-dimethylphenoxy)methyl]oxirane; bp. 85-90°C at 0.2 mm pressure (127); and

[(2-bromo-4-fluorophenoxy)methyl]oxirane; bp. 105°C at 66.5 Pa (128).

Example 16

20 To a stirred and cooled (temp. < 0°C) solution of 63 parts of 2-phenylcyclopropanemethanol in 378 parts of 1,1'-oxybisethane were added dropwise 63 parts of phosphor tribromide. The mixture was allowed to reach room temperature while stirring and the whole was further stirred for 30 minutes at room temperature. The reaction
25 mixture was poured onto 90 parts of ice water and the layers were separated. The organic layer was washed successively twice with water and once with a sodium hydrogen carbonate solution, dried, filtered and evaporated. The residue was distilled, yielding 67 parts (74.5%) of (1-bromo-3-butenyl)benzene; bp. 85-94°C at 0.4
30 mm pressure (129).

Example 17

A ballon was charged with 32 parts of 1-[(3-chloropropyl)-thio]-4-fluorobenzene, 50 parts of hydrogen peroxide and 112.5 parts of glacial acetic acid: exothermic reaction, the temperature rose to
35 about 60°C. The whole was stirred and refluxed for one hour and

further cooled while stirring for 3 hours. The cooled reaction mixture was then decomposed with 500 parts of water. The aqueous layer was separated and extracted once with 1,1'-oxybisethane. The combined organic layers were washed with water, dried over sodium sulfate and evaporated. The residue was recrystallized from 2,2'-oxybispropane, yielding, after cooling to -20°C , 18 parts of 1-[(3-chloropropyl)sulfonyl]-4-fluorobenzene (130).

Example 18

To a stirred and refluxing mixture of 56 parts of lithium aluminum hydride in 1000 parts of tetrahydrofuran was added dropwise a solution of 359 parts of 2-(2,5-dimethylphenoxy)propanoic acid in 1000 parts of tetrahydrofuran. Upon completion, stirring was continued at reflux temperature for 1.50 hours. The reaction mixture was cooled and decomposed by the successive additions of 55 parts of water, 45 parts of a 15% sodium hydroxide solution and 190 parts of water and the whole was stirred for 30 minutes. The inorganic material was filtered off after acidification with hydrochloric acid solution. The filtrate was dried, filtered and evaporated. The residue was distilled, yielding 73 parts of 2-(2,5-dimethylphenoxy)-1-propanol; bp. 150°C at 11 mm pressure (131).

Example 19

To a stirred and cooled mixture of 33.25 parts of 2-(phenylmethoxy)-1-propanol in 20 parts of pyridine was added dropwise a solution of 25.2 parts of methanesulfonyl chloride in 8 parts of pyridine at a temperature between 0° and 5°C . Upon completion, stirring was continued for 2 hours at room temperature. The reaction mixture was poured onto a mixture of crushed ice and ice water and the product was extracted three times with trichloromethane. The combined extracts were dried, filtered and evaporated. The oily residue was distilled, yielding 41 parts of 2-(phenylmethoxy)-1-propanol, methanesulfonate ester; bp. $151-152^{\circ}\text{C}$ at 0.03 mm pressure (132).

In a similar manner there were also prepared:

2-(3-methylphenoxy)-1-propanol, methanesulfonate ester; bp. 120°C at 0.0002 mm Hg pressure (133);

- 2-(2,6-dimethylphenoxy)-1-propanol, methanesulfonate ester; bp.
110°C at 0.0002 mm. pressure (134);
2-(3,5-dimethylphenoxy)-1-propanol, methanesulfonate ester; bp.
115°C at 0.0004 mm. pressure (135);
5 2-(2,6-dichlorophenoxy)-1-propanol, methanesulfonate ester; bp.
120°C at 0.002 mm. pressure (136);
3,4-dihydro-2H-1-benzopyran-2-methanol 4-methylbenzenesulfonate
ester; mp. 59.4°C (137).

Example 20

- 10 A mixture of 6.72 parts of 4-fluorophenol, 8.4 parts of potassium
carbonate and 200 parts of 4-methyl-2-pentanone was distilled
azeotropically to dry. 17.1 Parts of 4-(phenylmethoxy)cyclohexyl
methanesulfonate were added and the whole was stirred and refluxed
over week-end using a water separator. After cooling, water was
15 added and the layers were separated. The organic layer was washed
with a sodium hydroxide solution and with water, dried, filtered and
evaporated, yielding 17.7 parts (98.3%) of 1-fluoro-4-[[4-(phenyl-
methoxy)cyclohexyl]oxy]benzene as a residue (138).

- A mixture of 17.1 parts of 1-fluoro-4-[[4-(phenylmethoxy)cyclo-
20 hexyl]oxy]benzene and 160 parts of methanol was hydrogenated at
normal pressure and at room temperature with 2 parts of palladium-
on-charcoal catalyst 10%. After the calculated amount of hydrogen
was taken up, the catalyst was filtered off over Hyflo and the
filtrate was evaporated, yielding 8.9 parts (74.3%) of 4-(4-fluoro-
25 phenoxy)cyclohexanol as a residue (139).

- To a stirred and cooled mixture of 8.9 parts of 4-(4-fluoro-
phenoxy)cyclohexanol and 160 parts of 2-propanone were added drop-
wise 13 parts of a solution of 26.7 parts of chromium trioxide in
42.3 parts of concentrate sulfuric acid diluted with water till a
30 volume of 100 parts at a temperature below 5°C. Upon completion,
stirring was continued for 3 hours at room temperature. After the
addition of a few parts of methanol, the reaction mixture was poured
onto water. The product was extracted with dichloromethane. The
extract was dried, filtered and evaporated, yielding 8.4 parts
35 (96.1%) of 4-(4-fluorophenoxy)cyclohexanone as a residue (140).

Example 21

To a stirred and cooled (ice water bath) mixture of 72 parts of 2-ethenyl-3,4-dihydro-2H-1-benzopyran, 45 parts of sodium hydrogen carbonate and 2860 parts of dichloromethane were added quickly 102.4 parts of 3-chlorobenzenecarboperoxoic acid at about 10°C. The mixture was allowed to reach room temperature and stirring was continued for 4 days. The reaction mixture was filtered and the filtrate was washed successively with water, a saturate sodium sulfite solution, a 5% sodium hydroxide solution and again with water. The organic phase was dried, filtered and evaporated. The oily residue was purified by column chromatography over silica gel using a mixture of trichloromethane and petroleum ether (50:50 by volume) as eluent.

The first fraction (unreacted starting material) was collected and the eluent was evaporated. The residue was taken up in 1300 parts of dichloromethane and 30 parts of 3-chlorobenzenecarboperoxoic acid were added at about 10°C. After stirring for 8 hours at room temperature another 30 parts of 3-chlorobenzenecarboperoxoic acid were added and stirring was continued for 12 hours. The mixture was filtered and the filtrate was washed successively with water, a saturate sodium sulfite solution, a 5% sodium hydroxide solution and again with water. The organic phase was dried, filtered and evaporated. The oily residue was purified by column chromatography over silica gel using a mixture of trichloromethane and petroleum ether (50:50 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 14 parts of crude product. The second fraction (A+B) was collected and the eluent was evaporated, yielding 28.0 parts of crude product.

The combined crude fractions (resp. 14 and 28.0 parts) were separated by HPLC over silica gel using methylbenzene as eluent. The first fraction (A-isomer) was collected, yielding 12.5 parts of (A)-3,4-dihydro-2-oxiranyl-2H-1-benzopyran as a residue (141). The second fraction (B-isomer) was collected and the eluent was evaporated, yielding 14 parts of (B)-3,4-dihydro-2-oxiranyl-2H-1-benzopyran as a residue (142).

0184257

Sample 22

From a stirred mixture of 82 parts of a sodium methoxide solution and 315 parts of *N,N*-dimethylformamide was distilled off 160 parts of the solvent (the temperature rose to 130°C). After cooling to room temperature, there were added first 98 parts of trimethylsulfoxide. The whole was stirred first for 30 minutes at room temperature and then for 1 hour at 50°C. The mixture was cooled to room temperature and 30 parts of 4-(4-fluorophenoxy)-2-propanone were added. During a period of 30 minutes. Upon completion, stirring continued for 1 hour at room temperature and 2 hours at 50°C. After stirring overnight at room temperature, the reaction mixture was poured onto ice water. The product was extracted with 2,2'-oxybispropane. The organic layer was dried, filtered and evaporated, yielding 54 parts (84%) of 2-[(4-fluorophenoxy)methyl]-2-methyloxirane as a residue (143).

In a similar manner there was also prepared:

(S)-3,4-dihydro-2-oxiranyl-2H-1-benzopyran (144).

Sample 23

20 To a stirred solution of 2.7 parts of sodium in 50 parts of 2-methoxyethanol were added 13.2 parts of 4-fluorophenol. The whole was stirred for 15 minutes. A solution of 25.2 parts of (R)-2,2-dimethyl-4-methylene-1,3-dioxolane in 100 parts of 2-propanone was added. The mixture was stirred for 12 hours at reflux temperature. After cooling, the mixture was poured onto ice water. The product was extracted with methylbenzene. The extract was dried and evaporated, yielding 26 parts (97.3%) of (S)-4-[(4-fluorophenoxy)methyl]-2,2-dimethyl-1,3-dioxolane as a residue (145).

30 A mixture of 26 parts of (S)-4-[(4-fluorophenoxy)methyl]-2,2-dimethyl-1,3-dioxolane, 35 parts of a 10% aqueous acid solution and 80 parts of 2-propanone was stirred for 2 hours at reflux temperature. The mixture was stirred overnight at room temperature, diluted with 240 parts of ethanol and evaporated. The residue was dissolved in trichloromethane. The organic layer was washed twice

with water, dried, filtered and evaporated. The residue was crystallized from tetrachloromethane. The product was filtered off and dried, yielding 11 parts (50.4%) of (-)-(R)-3-(4-fluorophenoxy)-1,2-propanediol; $[\alpha]_D = -10.08^\circ$ (c = 0.5% in methanol) (146).


5 To a stirred mixture of 11 parts of (-)-(R)-3-(4-fluorophenoxy)-1,2-propanediol, 23.3 parts of pyridine and 240 parts of trichloromethane were added 12.3 parts of 4-methylbenzenesulfonyl chloride. The whole was stirred overnight at room temperature. It was washed successively with acid water, a sodium carbonate solution and water, dried, filtered and evaporated. The residue was purified
10 by column chromatography over silica gel using a mixture of trichloromethane and methanol (98:2 by volume) as eluent. The pure fractions were collected and the eluent was evaporated, yielding 13.5 parts (67.2%) of (+)-(S)-3-(4-fluorophenoxy)-1,2-propanediol,
15 α^1 -4-methylbenzenesulfonate; $[\alpha]_D = +11.11^\circ$ (c = 0.5% in trichloromethane) (147).

To a stirred mixture of 13.5 parts of (+)-(S)-3-(4-fluorophenoxy)-1,2-propanediol, α^1 -4-methylbenzenesulfonate and 45 parts of tetrahydrofuran were added 4.9 parts of potassium 1,1-dimethylethoxide. The whole was stirred for 30 minutes at room temperature.
20 It was filtered over Hyflo, washed with tetrahydrofuran and the

at 66.5 Pa; $[\alpha]_D = +5.27$ (c = 0.5% in dichloromethane); bp. 52-53°C at 66.5 Pa;
25 In a similar manner was also prepared:
(+)-(S)-[(4-fluorophenoxy)methyl]methane; bp. 52-53°C at 66.5 Pa;
 $[\alpha]_D = +5.27$ (c = 0.5% in dichloromethane) $[\alpha]_D = +9.1$ (c = 0.5% in dichloromethane)

B) Preparation of Final compoundsExample 24

A mixture of 4.4 parts of 1-chloro-3-(4-chlorobutoxy)benzene, 3.7 parts of N-methyl-N-(4-piperidiny1)-2-benzothiazolamine, 2.1 parts of sodium carbonate and 90 parts of N,N-dimethylformamide was stirred for 24 hours at 60°C. The reaction mixture was cooled and poured onto water. The product was extracted with methylbenzene. The extract was washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (Z)-2-butenedioate salt in 2-propanone. The salt was filtered off and dried, yielding 4 parts (49%) of N-[1-[4-(3-chlorophenoxy)butyl]-4-piperidiny1]-N-methyl-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 138.6°C (1).



No.	Ar	s	R	R ¹	Z	A	base salt	mp in °C °C
25	2 C ₆ H ₅	3	H	H	O	CH	base	123.2
3	C ₆ H ₅	3	H	CH ₃	O	CH	base	95.1
4	C ₆ H ₅	2	H	CH ₃	O	CH	1 l/2HNO ₃	126
5	C ₆ H ₅	2	H	H	O	CH	base	132
30	6 4-Cl-C ₆ H ₄	3	H	CH ₃	S	CH	base	111.9
7	C ₆ H ₅	3	H	CH ₃	S	CH	base	97.6
8	4-Cl-C ₆ H ₄	3	H	CH ₃	O	CH	base	87.5
9	3-F-C ₆ H ₄	3	H	CH ₃	S	CH	base	92.6
10	C ₆ H ₅	4	H	CH ₃	S	CH	*	154.4
35	11 4-F-C ₆ H ₄	4	H	CH ₃	S	CH	base	71.0

	12	4-Cl-C ₆ H ₄	4	H	CH ₃	S	CH	*	142.7
	13	C ₆ H ₅	2	H	CH ₃	S	CH	*	174.5
	14	4-CH ₃ -C ₆ H ₄	3	H	CH ₃	S	CH	base	90.0
5	15	4-Cl-C ₆ H ₄	4	H	CH ₃	O	CH	base	96.4
	16	4-F-C ₆ H ₄	4	H	CH ₃	O	CH	base	88.4
	17	4-Cl-C ₆ H ₄	2	H	CH ₃	S	CH	base	89.2
	18	2-F-C ₆ H ₄	3	H	CH ₃	S	CH	base	129.3
	19	4-CH ₃ O-C ₆ H ₄	3	H	CH ₃	O	CH	base	79.3
10	20	4-CH ₃ -C ₆ H ₄	3	H	CH ₃	O	CH	*	200.2
	21	4-1-C ₃ H ₇ -C ₆ H ₄	3	H	CH ₃	S	CH	*	180.4
	22	2,4-(Cl) ₂ -C ₆ H ₃	3	H	CH ₃	S	CH	*	179.3
	23	4-Br-C ₆ H ₄	3	H	CH ₃	S	CH	*	167.2
	24	4-CH ₃ O-C ₆ H ₄	3	H	CH ₃	S	CH	base	146.5
15	25	1-naphthalenyl	3	H	CH ₃	S	CH	*	154.9
	26	4-CH ₃ O-C ₆ H ₄	3	H	CH ₃	S	CH	base	156.0
	27	4-CH ₃ -C ₆ H ₄	2	H	CH ₃	S	CH	*	172.3
	28	4-CH ₃ -C ₆ H ₄	4	H	CH ₃	S	CH	*	155.5
	29	4-Cl,2-CH ₃ -C ₆ H ₄	3	H	CH ₃	S	CH	base	106.8
20	30	2-Cl-C ₆ H ₄	3	H	CH ₃	S	CH	base	111.4
	31	4-CH ₃ O-C ₆ H ₄	4	H	CH ₃	S	CH	base	89.4
	32	4-F-C ₆ H ₄	5	H	CH ₃	S	CH	*	143.5
	33	4-C ₆ H ₅ -C ₆ H ₄	3	H	CH ₃	S	CH	base	126.6
	34	2,6-Cl ₂ -C ₆ H ₃	3	H	CH ₃	S	CH	base	116.9
25	35	3,4,5-Cl ₃ -C ₆ H ₂	3	H	CH ₃	S	CH	*	181.3
	36	C ₆ H ₅	6	H	CH ₃	S	CH	**	127.6
	37	2-NH ₂ CO,4-F-C ₆ H ₃	3	H	CH ₃	S	CH	base	146.4
	38	C ₆ H ₅	4	H	CH ₃	O	CH	base	71.8
	39	2,6-(CH ₃) ₂ -C ₆ H ₃	3	H	CH ₃	S	CH	*	175.6
30	40	3-CH ₃ -C ₆ H ₄	3	H	CH ₃	S	CH	base	86.3
	41	2-CH ₃ -C ₆ H ₄	3	H	CH ₃	S	CH	base	93.2
	42	4-Br-C ₆ H ₄	4	H	CH ₃	S	CH	**	120.5
	43	4-CH ₃ O-C ₆ H ₄	5	H	CH ₃	S	CH	*	150.7
	44	2,4,5-Br ₃ -C ₆ H ₂	3	H	CH ₃	S	CH	*	164.2
35	45	4-CH ₃ -C ₆ H ₄	4	H	CH ₃	O	CH	base	85.4

[illegible]


	76	4-F-C ₆ H ₄	4	H	H	S	N	base	146.5
	77	3-F-C ₆ H ₄	3	H	H	S	N	base	155.8
	78	C ₆ H ₅	4	H	H	S	N	base	137.6
5	79	4-CH ₃ O-C ₆ H ₄	4	H	H	S	N	base	104.2
	80	3-Cl-C ₆ H ₄	4	H	H	S	N	base	110.7
	81	3-CH ₃ -C ₆ H ₄	3	H	H	S	N	base	163.9
	82	C ₆ H ₅	4	H	CH ₃	S	N	*	164.3
	83	3-F-C ₆ H ₄	3	H	CH ₃	S	N	*	144.9
10	84	3-Cl-C ₆ H ₄	4	H	CH ₃	S	N	*	148.3

* : (Z)-2-butenedioate (1:1)

** : (Z)-2-butenedioate (1:2)

*** : (E)-2-butenedioate (1:1)

15 In a similar manner there were also prepared:




No.	Ar	s	A ¹ -A ² -A ³ -A ⁴	base/ salt	mp in °C
20	85	3	CH=CH-C=CH OH	base	151.8
25	86	3	CH=CH-C=CH OH	base	135.5
	87	3	CH=C-CH=CH OH	2(COOH) ₂	176.3
	88	3	CH=C-CH=CH OH	****	212.2
30	89	3	CH=CCl-CH=CH	base	89.8
	90	4	CH=CCl-CH=CH	base	84.7
	91	4	CH=CCl-CH=CH	base	94.5
	92	3	CH=CCl-CH=CH	base	123.2

35

**** : (E)-2-butenedioate (2:1)

In a similar manner there were also prepared:



5	No.	L	R ¹	Z	A	base mp in	
						salt	°C
	93	n.butyl	H	S	N	base	170.5
10	94		H	S	N	base	157.9
	95	n.butyl	H	O	CH	base	139.2
	96	n.hexyl	H	S	N	base	152
	97	n.butyl	CH ₃	S	N	HCl	273.6 (dec.)
15	98	C ₆ H ₅ (CH ₂) ₂	H	O	CH	base	130.5
	99		H	O	CH	base	144.6
	100	n.hexyl	CH ₃	O	CH	base	54.8
20	101	4-CH ₃ O-C ₆ H ₄ -(CH ₂) ₂ -	CH ₃	O	CH	base	113.1
	102	C ₆ H ₅ CH=CH-CH ₂	CH ₃	O	CH	2HBr	247.2
	103	n.butyl	CH ₃	O	CH	2HBr	230.9
25	104		CH ₃	O	CH	2HBr	192.8
	105	4-CH ₃ O-C ₆ H ₄ -(CH ₂) ₂ -	H	O	CH	base	147.6
	106	n.hexyl	H	O	CH	2HNO ₃	141.5
	107	C ₆ H ₅ CH=CH-CH ₂ -	H	O	CH	base	123.4
30	108	C ₆ H ₅ -(CH ₂) ₅ -	CH ₃	S	CH	*	152.5
	109	n.undecyl	CH ₃	S	CH	*	163.5
	110	C ₆ H ₅ CH ₂ O-(CH ₂) ₃ -	CH ₃	S	CH	*	135.2
35	111	C ₆ H ₅ CH(OCH ₃)-	CH ₃	S	CH	*	150.3

	112	$\text{HO}-(\text{CH}_2)_3^-$	CH_3	S	CH	base	89.2
	113	$(4-\text{F}-\text{C}_6\text{H}_4)_2\text{CHO}$ $-(\text{CH}_2)_2^-$	CH_3	S	CH	$(\text{COOH})_2$	171.1
5	114	$4-\text{F}-\text{C}_6\text{H}_4-\text{SO}_2$ $-(\text{CH}_2)_3^-$	CH_3	S	CH	base	130.4

* : (Z)-2-butenedioate (1:1)

And following the same procedures and using the appropriate

10 starting materials there were also prepared:

1-butyl-N-(1-methylthiazolo[5,4-b]pyridin-2(1H)-ylidene)-4-piperidinamine dihydrochloride monohydrate; mp. 253.1°C (115);

N-(1-methylthiazolo[5,4-b]pyridin-2(1H)-ylidene)-1-(2-phenylethyl)-4-piperidinamine dihydrochloride monohydrate; mp. 251.7°C (116);

15 N-methyl-N-[1-(1-phenyl-3-butenyl)-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate (1:2) monohydrate; mp. 101.7°C (117);

N-[1-[4-(4-fluorophenoxy)butyl]-3-pyrrolidinyl]-N-methyl-2-benzothiazolamine dihydrochloride monohydrate; mp. 152.5°C (118);

20 N-[1-[4-(3-chlorophenoxy)butyl]-3-pyrrolidinyl]-N-methyl-2-benzothiazolamine ethanedioate (1:1); mp. 162.7°C (119);

N-methyl-N-[1-[3-(3-methylphenoxy)propyl]-3-pyrrolidinyl]-2-benzothiazolamine ethanedioate (1:1); mp. 108.4°C (120);

N-[1-[3-(3-fluorophenoxy)propyl]-3-pyrrolidinyl]-N-methyl-2-benzothiazolamine ethanedioate (1:1); mp. 137.0°C (121); and


25 N-methyl-N-[1-(4-phenoxybutyl)-3-pyrrolidinyl]-2-benzothiazolamine ethanedioate (1:1); mp. 150.2°C (122).

Example 25

A mixture of 5.04 parts of 1-(3-bromopropoxy)-4-methylbenzene, 7.54 parts of N-ethyl-N-(4-piperidinyl)-2-benzothiazolamine
30 (Z)-2-butenedioate (1:1), 5.3 parts of sodium carbonate, 0.1 parts of potassium iodide and 180 parts of N,N-dimethylformamide was stirred and heated overnight at 60°C. The reaction mixture was poured onto water and the product was extracted with methylbenzene. The extract was washed three times with water, dried, filtered and

evaporated. The residue was crystallized from 2-propanol, yielding 6 parts (75%) of *N*-ethyl-*N*-[1-[3-(4-methylphenoxy)propyl]-4-piperidinyl]-2-benzothiazolamine; mp. 101.0°C (123).

Following the same procedures and, if desired, after converting
5 the reaction product into the desired acid addition salt, there were also prepared:




10

No.	L'	s	R	R ¹	Z	base/salt	mp. in °C
15	124 C ₆ H ₅	2	H	H	S	base	118.1
	125 C ₆ H ₅	2	OH	H	S	base	140.1
						(cis-isomer)	
	126 2,3-dihydro-1,4-benzo-dioxin-2-yl	1	H	CH ₃	S	base	123.6
20	127 H	4	H	CH ₃	S	HCl	232.6
	128 H	4	H	H	S	base	116.8
	129 2,3-dihydro-1,4-benzo-dioxin-2-yl-	1	H	H	S	base	154.7
	130 4-F-C ₆ H ₄ -S-	3	H	CH ₃	S	*	158.0
25	131 4-F-C ₆ H ₄ -S-	3	H	CH ₃	O	base	87.5
	132 4-F-C ₆ H ₄ -S-	3	H	CH ₃	O	*	173.4
	133 C ₆ H ₅ -CH=CH-	1	H	CH ₃	S	base	101.3
	134 1H-benzimidazol-1-yl-	3	H	CH ₃	S	**	168.7
	135 4-F-C ₆ H ₄ -CH(CN)-	3	H	CH ₃	S	(COOH) ₂	190.1
30	136 HOCH ₂ -CH(OH)-	1	H	CH ₃	S	2HCl.H ₂ O	218.4
	137 C ₆ H ₅ -NH-	2	H	CH ₃	S	3HCl.2H ₂ O	204.7
	138 (C ₆ H ₅) ₂ CH-O-	2	H	CH ₃	S	(COOH) ₂ 1/2 H ₂ O	190.4
	139 C ₆ H ₅ CH(CH ₃)-O-	2	H	CH ₃	S	*	165.0
35	140 H	10	H	CH ₃	S	*	165.3
	141 5-phenyl-3-isoxazolyl	2	H	CH ₃	S	*	183.0

5	142 HO-	2 H	CH ₃	S *	175.0
	143 4-F-C ₆ H ₄ CH-	1 H	CH ₃	S *	180.5
	144 4-F-C ₆ H ₄ CH(OH)-	3 H	CH ₃	S **	145.5
	145 C ₆ H ₅ -CH=CH	1 H	C ₂ H ₅	S (COOH) ₂	199.6

In a similar manner there were also prepared:



No.	Ar	s	R ¹	Z base/salt	mp. in °C
15	146 4-F-C ₆ H ₄	3	CH ₃	S base	97.7
	147 4-F-C ₆ H ₄	3	CH ₃	O base	113.1
	148 2-naphthalenyl	3	CH ₃	S base	113.7
	149 2,6-Br ₂ -4-CH ₃ C ₆ H ₂	3	CH ₃	S *	169.7
	150 4-C ₂ H ₅ -C ₆ H ₄	5	CH ₃	S *	163.1
20	151 4-NO ₂ -C ₆ H ₄	3	CH ₃	S base	138.9
	152 2-naphthalenyl	5	CH ₃	S *	193.9
	153 3-CF ₃ -C ₆ H ₄	3	CH ₃	S *	149.4
	154 3-CH ₃ O-C ₆ H ₄	4	CH ₃	S *	125.8
	155 2-F-C ₆ H ₄	5	CH ₃	S *	165.8
25	156 4-C ₂ H ₅ -C ₆ H ₄	3	CH ₃	S *	176.6
	157 3-CH ₃ O-C ₆ H ₄	5	CH ₃	S *	157.3
	158 2,5-Cl ₂ -C ₆ H ₃	3	CH ₃	S *	171.7
	159 2-CH ₃ CO-4-F-C ₆ H ₃	3	CH ₃	S *	144.7
	160 3-F-C ₆ H ₄	5	CH ₃	S *	186.7
30	161 4-CH ₃ O-C ₆ H ₄	4	H	S base	95.4
	162 4-F-C ₆ H ₄	4	H	S base	125.8
	163 4-Cl-C ₆ H ₄	4	H	S base	149.8
	164 4-Cl-C ₆ H ₄	4	C ₂ H ₅	S *	136.5

165	4-CH ₃ O-C ₆ H ₄	4	C ₂ H ₅	S *	125.1
166	4-F-C ₆ H ₄	4	C ₂ H ₅	S (COOH) ₂	163.9
167	3-F-C ₆ H ₄	6	CH ₃	S ***	155.9

5

* : (Z)-2-butenedioate (1:1)

** : (Z)-2-butenedioate (1:2)

***: (E)-2-butenedioate (1:1)

In a similar manner there were also prepared:

- 10 N-[1-[3-(4-fluorophenoxy)-2-methylpropyl]-4-piperidiny]-N-methyl-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 143.6°C (168);
- 5-methoxy-N-[1-[4-(4-methoxyphenoxy)butyl]-4-piperidiny]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 98.2°C (169);
- N-(3-methyl-2(3H)-benzothiazolylidene)-1-[3-(3-methylphenoxy)propyl]-4-piperidinamine (Z)-2-butenedioate(1:2); mp. 176.0°C (170);
- 15 N-(3-methyl-2(3H)-benzothiazolylidene)-1-(4-phenoxybutyl)-4-piperidinamine (Z)-2-butenedioate(1:2); mp. 187.6°C (171);
- 1-[4-(3-chlorophenoxy)butyl]-N-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine (Z)-2-butenedioate(1:2); mp. 164.6°C (172);
- 20 1-[4-(4-fluorophenoxy)butyl]-N-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine (Z)-2-butenedioate(1:2); mp. 172.6°C (173);
- 1-[4-(4-methoxyphenoxy)butyl]-N-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine (Z)-2-butenedioate(1:2); mp. 166.6°C (174);
- N-(3-methyl-2(3H)-benzothiazolylidene)-1-(3-phenyl-2-propenyl)-4-piperidinamine (Z)-2-butenedioate (1:2); mp. 179.4°C (175);
- 25 5-bromo-N-[1-[4-(4-fluorophenoxy)butyl]-4-piperidiny]-6-methoxy-N-methyl-2-benzothiazolamine; mp. 124.6°C (176);
- N-[1-[3-(3-fluorophenoxy)propyl]-4-piperidiny]-5-methoxy-N-methyl-2-benzothiazolamine (Z)-2-butenedioate(1:1); mp. 126.3°C (177).

30 Example 26

A mixture of 2.3 parts of [2-(4-methoxyphenyl)ethyl] methanesulfonate, 4 parts of N-(4-piperidiny)-2-benzothiazolamine dihydrobromide, 5.3 parts of sodium carbonate, 0.1 parts of potassium iodide and 90 parts of N,N-dimethylformamide was stirred overnight at 70°C. The

35 reaction mixture was poured onto water and the product was extracted

with methylbenzene. The extract was washed with water, dried, filtered and evaporated. The residue was crystallized from a mixture of 2-propanol and 2,2'-oxybispropane. The product was filtered off and dried, yielding 1.5 parts (41%) of N-[1-[2-(4-methoxyphenyl)-ethyl]-4-piperidinyl]-2-benzothiazolamine; mp. 142.5°C (178).

In a similar manner there were also prepared:


N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 97.6°C (179); and

N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]thiazolo[5,4-b]pyridin-2-amine; mp. 158.9°C (180).

Example 27

A mixture of 4.3 parts of 1-(4-chlorobutoxy)-4-methoxybenzene, 6.6 parts of N-(phenylmethyl)-N-(4-piperidinyl)-2-benzothiazolamine (Z)-2-butenedioate(1:1), 5.3 parts of sodium carbonate, 0.1 parts of potassium iodide and 90 parts of N,N-dimethylacetamide was stirred overnight at 90°C. The reaction mixture was poured onto water. The product was extracted with methylbenzene. The extract was washed with water, dried, filtered and evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 5.6 parts (60%) of N-[1-[4-(4-methoxyphenoxy)butyl]-4-piperidinyl]-N-(phenylmethyl)-2-benzothiazolamine (E)-2-butenedioate(1:1); mp. 178.2°C (181).

In a similar manner there were also prepared:




No.	L'	s	Z	base/salt	mp. in °C
182	H	12	S	*	161.4
183	3-oxo-1,2,4-triazolo-[4,5-a]pyridin-2-yl	3	S	*	185.0
184	4-F-C ₆ H ₄ -CH(CN)-	3	O	1 1/2(COOH) ₂	139.5

5	185	$(C_6H_5)_2CH-O-$	2	O	*	175.3
	186	HO	2	O	*	172.6
	187	C_6H_5-NH-	3	S	$2 \frac{1}{2}(COOH)_2$	177.1
	188	2,3-dihydro-1,4-benzo- dioxin-2-yl	2	O	*	182.1
	189	C_6H_5-NH-	2	O	*	152.6
10	190	2,3-dihydro-1,4-benzo- dioxin-2-yl	2	S	*	168.6

*: (Z)-2-butenedioate (1:1)

In a similar manner there were also prepared:



No.	Ar	s	R ¹	Z	base/ salt	mp. in °C
20	191 3-Cl-C ₆ H ₄ -	5	CH ₃	S	*	177.7
	192 4-CH ₃ -C ₆ H ₄ -	5	CH ₃	S	*	165.1
	193 2-F-C ₆ H ₄ -	3	CH ₃	O	*	208.0
	194 3,5-Cl ₂ -C ₆ H ₃ -	3	CH ₃	S	*	178.1
25	195 3-F-C ₆ H ₄ -	3	CH ₃	O	*	180.6
	196 3-F-C ₆ H ₄ -	4	CH ₃	S	*	143.8
	197 3-F-C ₆ H ₄ -	4	CH ₃	O	*	158.8
	198 C ₆ H ₅ -	4	C ₆ H ₅ CH ₂	S	*	129.9
	199 4-CH ₃ -C ₆ H ₄ -	3	C ₆ H ₅ CH ₂	S	**	198.4
30	200 4-Cl-C ₆ H ₄ -	4	C ₆ H ₅ CH ₂	S	*	149.5
	201 3-F-C ₆ H ₄ -	3	C ₆ H ₅ CH ₂	S	*	145.1
	202 4-F-C ₆ H ₄ -	4	C ₆ H ₅ CH ₂	S	*	153.6
	203 4-NO ₂ -C ₆ H ₄ -	4	CH ₃	S	*	169.7

35 *: (Z)-2-butenedioate (1:1)

** : (E)-2-butenedioate (1:1)


In a similar manner there was also prepared:

1-[3-(3-fluorophenoxy)propyl]-N-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine (2)-2-butenedioate(1:2); mp. 173.4°C (204).

Example 28


- 5 A mixture of 5.5 parts of cis-N-(3-methoxy-4-piperidinyl)-N-methyl-2-benzothiazolamine ethanedioate(1:1), 10.6 parts of sodium carbonate, 0.1 parts of potassium iodide and 200 parts of 4-methyl-2-pentanone was stirred and refluxed for 30 minutes. After cooling to 60°C, 4.4 parts of 1-chloro-4-(4-chlorobutoxy)benzene were added.
- 10 The whole was stirred and refluxed overnight. After cooling, water was added and the layers were separated. The organic layer was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (98:2 by volume) as eluent. The pure fractions were
- 15 collected and the eluent was evaporated. The residue was converted into the ethanedioate salt in 2-propanol. The salt was filtered off and dried, yielding 6 parts (73%) of cis-N-[1-[4-(4-chlorophenoxy)-butyl]-3-methoxy-4-piperidinyl]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 210.4°C (205).

- 20 In a similar manner there were also prepared:



25	No.	L'	s	R ¹	Z	A ¹ =A ² -A ³ =A ⁴	base salt	mp in °C
	206	3-oxo-1,2,4-triazolo- [4,3-a]pyridin-2-yl	3	CH ₃	S	N=CH-CH=CH	2HCl 1/2H ₂ O	235.5
	207	cyclohexyloxy	3	CH ₃	S	CH=CH-CH=CH	*	160.2
30	208	4-F-C ₆ H ₄ -CH=CH-	2	CH ₃	S	CH=CH-CH=CH	*	145.0
	209	3,4-dihydro-2H-1- benzopyran-2-yl	1	CH ₃	O	CH=CH-CH=CH	(COOH) ₂	212.1
	210	3,4-dihydro-2H-1- benzopyran-2-yl	1	CH ₃	S	CH=CH-CH=CH	base	129.8
35	211	C ₆ H ₅ -CH=CH-	1	H	S	CH=CH-CH=CH	**	177.8

```
* : (Z)-2-butenedioate (1:1)  ** : (Z)-2-butenedioate (1:2)
*** : (E)-2-butenedioate (1:1)
```



	No. Ar	S R	R ¹	Z	A ¹ =A ² -A ³ =A ⁴	cis base mp in /trans salt °C
25	225 2-CH ₃ OCO-C ₆ H ₄	3 H	CH ₃	S	CH=CH-CH=CH - *	177.0
	226 4-F-C ₆ H ₄	6 H	CH ₃	S	CH=CH-CH=CH - *	140.0
	227 C ₆ H ₅	5 H	CH ₃	S	CH=CH-CH=CH - *	179.9
	228 2-n.C ₃ H ₇ C ₆ H ₄	3 H	CH ₃	S	CH=CH-CH=CH - *	164.8
30	229 3-F-C ₆ H ₄	3 OCH ₃	CH ₃	S	CH=CH-CH=CH cis ***	210.5
	230 3-CH ₃ -C ₆ H ₄	3 H	CH ₃	S	CH=CF-CH=CH - base	90.6
	231 3-F-C ₆ H ₄	3 H	CH ₃	S	CH=C(CH ₃)-CH=CH - *	140.7
35	232 3-CH ₃ -C ₆ H ₄	3 H	CH ₃	S	CH=C(CH ₃)-CH=CH - *	165.6

	233	C_6H_5	4	H	CH_3	S	$CH=C-CH=CH$ $ $ CH_3	- *	141.4
	234	$4-CH_3O-C_6H_4$	4	H	CH_3	S	$CH=CF-CH=CH$	- base	90.7
5	235	$3-F-C_6H_4$	3	H	CH_3	S	$CH=CF-CH=CH$	- *	171.1
	236	C_6H_5	4	H	CH_3	S	$CH=CF-CH=CH$	- *	142.7
	237	$4-CH_3O-C_6H_4$	4	H	CH_3	S	$CH=C-CH=CH$ $ $ CH_3	- *	123.8
	238	$3-F-C_6H_4$	3	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- *	160.1
10	239	$4-F-C_6H_4$	4	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- *	161.1
	240	$4-Cl-C_6H_4$	4	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- *	149.7
	241	$4-CH_3-C_6H_4$	3	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- base	99.2
	242	$3-F-C_6H_4$	3	H	CH_3	S	$CH=CH-CCl=CH$	- *	153.8
	243	$4-CH_3O-C_6H_4$	4	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- *	158.7
15	244	C_6H_5	4	H	$1.C_3H_7$	S	$CH=CH-CH=CH$	- *	136.2
	245	$3-F-C_6H_4$	3	H	CH_3	S	$CCl=CH-CH=CH$	- *	181.2
	246	$3-CH_3-C_6H_4$	3	H	CH_3	S	$CH=CH-C-CH$ $ $ OCH_3	- **	161.2
	247	C_6H_5	4	OCH_3	CH_3	S	$CH=CH-CH=CH$	cis ****	198.3
20	248	$3-CH_3-C_6H_4$	3	OCH_3	CH_3	S	$CH=CH-CH=CH$	cis ****	187.2
	249	$4-CH_3O-C_6H_4$	4	OCH_3	CH_3	S	$CH=CH-CH=CH$	cis ****	214.6
	250	$3-CH_3-C_6H_4$	3	H	CH_3	S	$CH=CH-CCl=CH$	- *	151.0
	251	$3-CH_3-C_6H_4$	3	H	CH_3	S	$CCl=CH-CH=CH$	- *	178.4
	252	$4-CH_3O-C_6H_4$	4	H	CH_3	S	$CH=CH-CF=CH$	- *	157.2
25	253	$4-CH_3O-C_6H_4$	4	H	CH_3	S	$CCl=CH-CH=CH$	- *	171.1
	254	$3-CH_3-C_6H_4$	3	H	CH_3	S	$CH=CH-CF=CH$	- *	163.8
	255	C_6H_5	4	H	CH_3	S	$CH=CH-CF=CH$	- *	185.1
	256	$3-F-C_6H_4$	3	H	CH_3	S	$CH=CH-CF=CH$	- *	167.9
	257	C_6H_5	4	H	CH_3	S	$CH=CH-CCl=CH$	- *	200.4
30	258	$4-CH_3O-C_6H_4$	4	H	CH_3	S	$CH=CH-CCl=CH$	- *	143.1
	259	C_6H_5	4	H	CH_3	S	$CCl=CH-CH=CH$	- *	176.8
	260	$4-Cl-C_6H_4$	4	H	H	O	$CH=CH-CH=CH$	- 2HCl	251.2
	261	$4-F-C_6H_4$	4	H	H	O	$CH=CH-CH=CH$	- 2HCl	239.8
	262	$4-CH_3-C_6H_4$	3	H	H	O	$CH=CH-CH=CH$	- 2HCl	246.6

5	263	4-CH ₃ O-C ₆ H ₄	4	H	H	O	CH=CH-CH=CH	-	2HCl	237.3
	264	3-CH ₃ -C ₆ H ₄	3	H	H	O	CH=CH-CH=CH	-	2HCl.	219.2
									H ₂ O	
	265	3-Cl-C ₆ H ₄	4	H	H	O	CH=CH-CH=CH	-	2HCl	217.3
	266	C ₆ H ₅	4	H	H	O	CH=CH-CH=CH	-	2HCl.	231.7
10									1/2H ₂ O	
	267	3-F-C ₆ H ₄	3	H	H	O	CH=CH-CH=CH	-	2HCl.	224.8
									H ₂ O	
	268	C ₆ H ₅	4	OH	H	S	CH=CH-CH=CH	cis base		158.2
	269	3-CH ₃ -C ₆ H ₄	3	OH	H	S	CH=CH-CH=CH	cis base		146.9
	270	4-Cl-C ₆ H ₄	4	OH	H	S	CH=CH-CH=CH	cis base		179.7
	271	3-F-C ₆ H ₄	3	OH	H	S	CH=CH-CH=CH	cis base		165.8

- 15 * : (Z)-2-butenedioate(1:1)
 ** : (E)-2-butenedioate(1:1)
 *** : ethanedioate (1:1) hemihydrate
 **** : ethanedioate (1:1)

In a similar manner there were also prepared:

- 20 N-methyl-N-[1-[2-(phenylmethoxy)propyl]-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 150.5°C (272);
N-methyl-N-[1-(2-methyl-3-phenoxypropyl)-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 177.0°C (273);
N-methyl-N-[1-[2-(3-methylphenoxy)propyl]-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 168.6°C (274);
 25 N-[1-[2-(2,6-dimethylphenoxy)propyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 130.4°C (275);
N-[1-[2-(2,6-dichlorophenoxy)propyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 108.0°C (276);
 30 N-[1-[2-(3,5-dimethylphenoxy)propyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 172°C (277);
N-(2-benzothiazolyl)-8-[3-(3-fluorophenoxy)propyl]-N-methyl-8-azabicyclo[3.2.1]octan-3-amine (E)-2-butenedioate(1:1); mp. 171.3°C (278);
 (E)-N-methyl-N-[8-(3-phenyl-2-propenyl)-8-azabicyclo[3.2.1]oct-3-yl]-
 35 2-benzothiazolamine ethanedioate(1:1); mp. 216.0°C (279);

- N-(2-benzothiazolyl)-N-methyl-8-(4-phenoxybutyl)-8-azabicyclo[3.2.1]-octan-3-amine (E)-2-butenedioate(1:1); mp. 158.0°C (280);
- cis-N-methyl-N-[3-methyl-1-(3-phenyl-2-propenyl)-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate(1:1); mp. 173.7°C (281);
- 5 cis-N-[3-methoxy-1-(3-phenyl-2-propenyl)-4-piperidinyl]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 230.2°C (282);
- N-(2-benzothiazolyl)-N-methyl-8-[3-(3-methylphenoxy)propyl]-8-azabicyclo[3.2.1]octan-3-amine ethanedioate(1:1); mp. 183.3°C (283);
- N-methyl-N-[1-(4-phenoxybutyl)-3-piperidinyl]-2-benzothiazolamine
- 10 ethanedioate(1:1); mp. 152.5°C (284);
- N-[1-[3-(3-fluorophenoxy)propyl]-3-piperidinyl]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 164.5°C (285);
- N-methyl-N-[1-(3-phenyl-2-propenyl)-3-piperidinyl]-2-benzothiazolamine ethanedioate(1:1); mp. 202.8°C (286);
- 15 N-methyl-N-[1-[3-(3-methylphenoxy)propyl]-3-piperidinyl]-2-benzothiazolamine ethanedioate(1:1); mp. 151.3°C (287);
- N-[1-[4-(4-methoxyphenoxy)butyl]-3-piperidinyl]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 152.3°C (288);
- N-[1-[4-(4-fluorophenoxy)butyl]-3-piperidinyl]-N-methyl-2-benzothiazolamine ethanedioate(1:1); mp. 163.2°C (289);
- 20 N-[1-[3-(4-fluorophenoxy)-1-methylpropyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 115.0°C (290);
- N-[1-[4-(3-chlorophenoxy)butyl]-3-piperidinyl]-N-methyl-2-benzothiazolamine (E)-2-butenedioate(1:1); mp. 138.6°C (291);
- 25 N-methyl-N-[1-(3-phenyl-2-propenyl)-3-pyrrolidinyl]-2-benzothiazolamine 4-methylbenzenesulfonate(1:2); mp. 198.8°C (292); and
- cis-4-(2-benzothiazolylamino)-1-(3-phenyl-2-propenyl)-3-piperidinol; mp. 198.3°C (293).

Example 29

- 30 A mixture of 3 parts of 1-bromo-3-methylbutane, 3.7 parts of N-methyl-N-(4-piperidinyl)-1-benzothiazolamine, 4.2 parts of potassium carbonate and 80 parts of 2-propanone was stirred and refluxed overnight. The reaction mixture was poured onto water and the product was extracted with methylbenzene. The extract was dried,
- 35 filtered and evaporated. The residue was converted into the (Z)-2-

butenedioate salt in 2-propanol. The salt was filtered off and dried, yielding 5 parts (77%) of N-methyl-N-[1-(3-methylbutyl)-4-piperidinyl]-2-benzothiazolamine (Z)-2-butenedioate (1:1); mp. 197.2°C (294).

5 Example 30

A mixture of 3 parts of (3-chloro-1-propenyl)benzene, 2.6 parts of 2-[methyl(4-piperidinyl)amino]-5-benzothiazolol, 3.5 parts of N,N-diethylethanamine and 90 parts of N,N-dimethylformamide was stirred and heated for 24 hours at 60°C. The reaction mixture was
10 cooled, poured onto water and the product was extracted with methylbenzene. The extract was washed with water, dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The pure fractions were collected and
15 the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 1 part (20%) of 2-[methyl[1-(3-phenyl-2-propenyl)-4-piperidinyl]amino]-5-benzothiazolol (E)-2-butenedioate(1:1); mp. 228.4°C (295).


20 Example 31

To a stirred mixture of 1.1 parts of a sodium hydride dispersion 50% and 45 parts of N,N-dimethylformamide were added 3.1 parts of 4-[(2-benzothiazolyl)methylamino]-1-piperidinepropanol and the whole was stirred and heated for 2 hours at about 60°C. After cooling to
25 about 15°C, 1.65 parts of 4-chloropyridine hydrochloride were added and stirring was continued overnight at 60°C. The reaction mixture was cooled and the solvent was evaporated. The residue was taken up in water and extracted with methylbenzene. The extract was dried, filtered and evaporated. The residue was purified by filtration over
30 silica gel using a mixture of trichloromethane and methanol (98:2 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 2,2'-oxybispropane, yielding, after drying, 1.5 parts (38%) of N-methyl-N-[1-[3-(4-pyridinyloxy)propyl]-4-piperidinyl]-2-benzothiazolamine
35 monohydrate; mp. 96.3°C (296).

Example 32

A mixture of 4.75 parts of cyclopentanone, 4.1 parts of N-(1-methylethyl)-N-(4-piperidinyl)-2-benzothiazolamine and 200 parts of methanol was hydrogenated at normal pressure and at room temperature with 2 parts of palladium-on-charcoal catalyst 10%. After the calculated amount of hydrogen was taken up, the catalyst was filtered off over Hyflo and the filtrate was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (98.5:1.5 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (Z)-2-butenedioate salt in 2-propanol. The salt was filtered off and dried, yielding 3.9 parts (56.6%) of N-(1-cyclopentyl-4-piperidinyl)-N-(1-methylethyl)-2-benzothiazolamine (Z)-2-butenedioate(1:1); mp. 178.5°C (297).

In a similar manner there were also prepared:



20

No. L	R ¹	Z	A ¹ =A ² -A ³ =A ⁴	base salt	mp in °C
25 298 CH ₃	H	S	CH=CH-CH=CH	2HCl. 1/2H ₂ O	283.1
299 CH ₃	H	S	N=CH-CH=CH	base	170.6
300 CH ₃	CH ₃	S	N=CH-CH=CH	2HCl	266.5
301 cyclohexyl	H	O	CH=CH-CH=CH	base	137.4
30 302 CH ₃	H	O	CH=CH-CH=CH	base	132.1
303 C ₂ H ₅	CH ₃	O	CH=CH-CH=CH	2HBr.H ₂ O	240.9
304 CH ₃	CH ₃	O	CH=CH-CH=CH	2HNO ₃	142.9
305 1.C ₃ H ₇	CH ₃	S	CH=CH-CH=CH	2HCl.H ₂ O	265.9
306 cyclohexyl	CH ₃	S	CH=CH-CH=CH	2HCl	278.8

35

	307 cyclopentyl	CH ₃	S	CH=CH-CH=CH	2HCl	268.8
	308 cyclopentyl	CH ₃	O	CH=CH-CH=CH	2HCl.	276.2-
					2H ₂ O	280.8(dec)
5	309 cyclohexyl	CH ₃	O	CH=CH-CH=CH	2HCl.H ₂ O	300
	310 cyclopropylmethyl	CH ₃	S	CH=CH-CH=CH	2HCl	236.4
	311 cyclopentyl	H	S	CH=CH-CH=CH	base	172.7
	312 cyclopentyl	C ₂ H ₅	S	CH=CH-CH=CH	*	145.1
	313 cyclopentyl	n.C ₄ H ₉	S	CH=CH-CH=CH	**	181.6
10	314 cyclopentyl	C ₆ H ₅ CH ₂	S	CH=CH-CH=CH	*	202.1
	315 cyclopropylmethyl	CH ₃	O	CH=CH-CH=CH	*	184.9

* : (Z)-2-butenedioate (1:1)

** : (E)-2-butenedioate (1:1)

15 In a similar manner there were also prepared:

cis-4-[4-[(2-benzothiazolyl)methylamino]-1-piperidinyl]-1-(4-fluorophenyl)cyclohexanecarbonitrile; mp. 184.7°C (316);

(cis+trans)-N-[1-[4-(4-fluorophenoxy)cyclohexyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 198.4°C (317);

20 cis-4-[4-[(2-benzoxazolyl)methylamino]-1-piperidinyl]-1-(4-fluorophenyl)cyclohexanecarbonitrile; mp. 199.6°C (318); and


N-[1-(1H-indol-3-ylmethyl)-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 178.3°C (319).

Example 33

25 A mixture of 2.5 parts of [(4-fluorophenoxy)methyl]oxirane, 3.7 parts of N-methyl-N-(4-piperidinyl)-2-benzothiazolamine, 45 parts of methylbenzene and 40 parts of methanol was stirred and refluxed for 24 hours. The reaction mixture was evaporated. The residue was

30 filtered off and dried, yielding 6 parts (85%) of 4-[(2-benzothiazolyl)methylamino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol monohydrochloride; mp. 168.4-178.8°C (320).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:




	No.	Ar	R ¹	Z base/ salt	mp. in °C
5	321	C ₆ H ₅	H	O base	119.8
	322	C ₆ H ₅	CH ₃	O HBr	231.6
	323	4-F-C ₆ H ₄	CH ₃	O HCl	233.5
10	324	4-Cl-C ₆ H ₄	CH ₃	O HCl	237.5
	325	2-C ₂ H ₅ O-C ₆ H ₄	CH ₃	O 2HCl	203.3
	326	2-propenyloxyphenyl	CH ₃	O HCl	144.6
	327	4-CH ₃ CONH-C ₆ H ₄	CH ₃	O 2HCl	236.0
	328	4-CH ₃ CONH-C ₆ H ₄	CH ₃	S 2HCl	169.6-182.2
15	329	1-naphthalenyl	CH ₃	S (COOH) ₂	198.1-199.9
	330	2-propenyloxyphenyl	CH ₃	S (COOH) ₂	120.3-129.5
	331	4-Cl-C ₆ H ₄	CH ₃	S 2HCl	206.0-225.6
	332	1-naphthalenyl	CH ₃	O (COOH) ₂	232.1-233.1
	333	2-acetylphenyl	CH ₃	O (COOH) ₂	203.2-203.6
20	334	2-acetylphenyl	CH ₃	S (COOH) ₂	222.8-223.1
	335	2-C ₂ H ₅ O-C ₆ H ₄	CH ₃	S (COOH) ₂	112.9-118.6
	336	2-Cl-C ₆ H ₄	CH ₃	H ₂ O S (COOH) ₂ 1/2H ₂ O	180.9-182.8
25	337	2-Cl-C ₆ H ₄	CH ₃	O (COOH) ₂ 1/2H ₂ O	172.6-182.1
	338	C ₆ H ₅	CH ₃	S (COOH) ₂	166.0-169.6
	339	2-CN-C ₆ H ₄	CH ₃	O (COOH) ₂	173.9-175.6
	340	2-CN-C ₆ H ₄	CH ₃	S *	150.5
30	341	3-Cl-C ₆ H ₄	CH ₃	S base	129.0
	342	2,4-Cl ₂ -C ₆ H ₃	CH ₃	O *	187.7
	343	2,4-Cl ₂ -C ₆ H ₃	CH ₃	S *	167.8

	344	3-CF ₃ -C ₆ H ₄	CH ₃	S	2HCl	220.9
	345	2,3-dihydro-1H-inden-5-yl	CH ₃	S	base	95.9
	346	2-Br-C ₆ H ₄	CH ₃	S	*	167.0
5	347	6-Br-2-napthalenyl	CH ₃	S	*	192.9
	348	4-Cl,3,5-(CH ₃) ₂ -C ₆ H ₂	CH ₃	S	base	131.7
	349	4-(4-methylbenzoyl)phenyl	CH ₃	S	base	174.2
	350	2,4,6-(Cl ₂) ₃ -C ₆ H ₂	CH ₃	S	base	126.4
	351	2,6-(CH ₃) ₂ -C ₆ H ₂	CH ₃	S	base	146.8
10	352	3-CH ₃ O-C ₆ H ₄	CH ₃	S	**	118.5
	353	2-n.C ₃ H ₇ -CO-C ₆ H ₄	CH ₃	S	(COOH) ₂	159.0
	354	3-Cl-C ₆ H ₄	CH ₃	O	*	123.7
	355	4-Cl-C ₆ H ₄	H	S	base	164.8
	356	4-F-C ₆ H ₄	H	S	base	162.8
15	357	4-Cl-C ₆ H ₄	C ₂ H ₅	S	base	103.1
	358	4-F-C ₆ H ₄	C ₂ H ₅	S	***	205.7
	359	4-Cl,2-CH ₃ -C ₆ H ₃	CH ₃	S	base	115.5
	360	2-F-C ₆ H ₄	CH ₃	S	base	100.4
	361	2-CH ₃ CO,4-F-C ₆ H ₃	CH ₃	S	base	108.7
20	362	4-F-C ₆ H ₄	CH ₃	S	***	169.5
	363	4-CH ₃ CO-C ₆ H ₄	CH ₃	S	base	98.9
	364	4-F-C ₆ H ₄	1.C ₃ H ₇	S	*	161.8
	365	4-F-C ₆ H ₄	CH ₃	S	base	101.7
	366	4-Cl-C ₆ H ₄	1.C ₃ H ₇	S	*	169.8
25	367	4-CN-C ₆ H ₄	CH ₃	S	base	122.1
	368	C ₆ H ₅	CH ₃	S	*	135.6
	369	4-F-C ₆ H ₄	CH ₃	S	2HCl	209.4
	370	4-NO ₂ -C ₆ H ₄	CH ₃	S	base	140.1
	371	4-F-C ₆ H ₄	H	O	2HCl	227.9
30	372	3-Cl-C ₆ H ₄	H	O	2HCl	213.2
	373	3,4,5-(CH ₃ O) ₃ -C ₆ H ₂	CH ₃	S	*	176.9
	374	4-CH ₃ O-C ₆ H ₄	CH ₃	S	base	129.6
	375	3,5-(CH ₃ O) ₂ -C ₆ H ₃	CH ₃	S	*	145.2
35	376	1,3-benzodioxol-5-yl	CH ₃	S	***	181.7

* : (Z)-2-butenedioate (1:1); ** : (Z)-2-butenedioate (1:2)

*** : (E)-2-butenedioate (2:1)

In a similar manner there were also prepared:



No.	Ar	R	R ¹	A ¹ =A ² -A ³ =A ⁴	cis/ base trans salt	mp in °C
10						
	377 C ₆ H ₅	CH ₃ O	CH ₃	CH=CH-CH=CH	cis *	163.6
	378 4-F-C ₆ H ₄	H	CH ₃	CH=CF-CH=CH	- base	101.5
	379 C ₆ H ₅	H	CH ₃	CH=CF-CH=CH	- base	130.6
	380 4-F-C ₆ H ₄	CH ₃ O	CH ₃	CH=CH-CH=CH	cis *	178.9
15	381 4-F-C ₆ H ₄	CH ₃	CH ₃	CH=CH-CH=CH	cis *	150.0
	382 C ₆ H ₅	CH ₃	CH ₃	CH=CH-CH=CH	cis *	165.1
	383 4-F-C ₆ H ₄	H	CH ₃	CH=CH-CF=CH	- base	118.1
	384 4-F-C ₆ H ₄	H	CH ₃	CCl=CH-CH=CH	- *	154.0
	385 4-F-C ₆ H ₄	H	CH ₃	CH=CH-CCl=CH	- *	144.6
20	386 4-F-C ₆ H ₄	H	H	N=CH-CH=CH	- base	168.2
	387 C ₆ H ₅	H	CH ₃	CH=CCl-CH=CH	- base	177.4
	388 4-F-C ₆ H ₄	H	CH ₃	CH=CCl-CH=CH	- base	172.7
	389 4-F-C ₆ H ₄	H	CH ₃	CH=CH-C=CH OCH ₃	- **	176.1
25	390 C ₆ H ₅	H	CH ₃	CCl=CH-CH=CH	- *	174.8
	391 C ₆ H ₅	H	CH ₃	CH=CH-CCl=CH	- *	171.1
	392 4-F-C ₆ H ₄	H	CH ₃	N=CH-CH=CH	- base	103.6
	393 3-Cl-C ₆ H ₄	H	CH ₃	N=CH-CH=CH	- base	119.9

30 * : (Z)-2-butenedioate (1:1)

** : (E)-2-butenedioate (1:1)

In a similar manner there were also prepared:

4-[(2-benzothiazolyl)methylamino]-α-(4-methoxyphenyl)-1-piperidine-ethanol; mp. 168.1°C (394);

35 4-[(2-benzothiazolyl)methylamino]-α-(4-fluorophenyl)-1-piperidine-

- ethanol; mp. 178.5°C (395);
- 3-[(2-benzothiazolyl)methylamino]- α -(phenoxymethyl)-8-azabicyclo[3.2.1]octane-8-ethanol ethanedioate(1:1); mp. 188.6°C (396);
- 3-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol ethanedioate (1:1); mp. 182.7°C (397);
- 3-[(2-benzothiazolyl)methylamino]- α -[(3-chlorophenoxy)methyl]-1-piperidineethanol ethanedioate (1:1); mp. 172.9°C (398);
- 3-[(2-benzothiazolyl)methylamino]- α -(phenoxymethyl)-1-piperidineethanol ethanedioate (1:1); mp. 180.2°C (399);
- 3-[(2-benzothiazolyl)methylamino]- α -[(3-chlorophenoxy)methyl]-1-pyrrolidineethanol ethanedioate(1:1); mp. 158.5°C (400);
- α -[(3-chlorophenoxy)methyl]-4-(thiazolo[5,4-b]pyridin-2-ylamino)-1-piperidineethanol; mp. 127.4°C (401);
- (A)-4-(2-benzoxazolylamino)- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-1-piperidineethanol dihydrochloride; mp. 229.8°C (402);
- (B)-4-(2-benzoxazolylamino)- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-1-piperidineethanol dihydrochloride.hemihydrate; mp. 245.3°C (403);
- 4-[(2-benzoxazolyl)methylamino]- α -(4-methoxyphenyl)-1-piperidineethanol dihydrochloride; mp. 231.1°C (404);
- 3-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-pyrrolidineethanol 4-methylbenzenesulfonate(1:2); mp. 199.8°C (405);
- α -[(4-fluorophenoxy)methyl]-4-[(1-methylthiazolo[5,4-b]pyridin-2-yliden)amino]-1-piperidineethanol; mp. 150.2°C (406);
- 4-[(3-methyl-2(3H)-benzothiazolyliden)amino]- α -(phenoxymethyl)-1-piperidineethanol (Z)-2-butenedioate(1:2); mp. 152.8°C (407);
- 4-[(2-benzothiazolyl)methylamino]- α -[[2,4-dichlorophenyl)methoxy]-methyl]-1-piperidineethanol ethanedioate(1:1); mp. 158.1°C (408);
- α -[(4-fluorophenoxy)methyl]-4-[(oxazolo[4,5-b]pyridin-2-yl)amino]-1-piperidineethanol (E)-2-butenedioate(1:1); mp. 195.4°C (409);
- (+)-(R)-4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol dihydrochloride mp. 179.2°C, $[\alpha] = +12.49^\circ$ (c = 0.5% in methanol) (410); and
- (-)-(S)-4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol dihydrochloride; mp. 175.7°C, $[\alpha] = -11.37^\circ$ (c = 0.5% in methanol) (411).

In a similar manner there is also prepared:

α -[(4-fluorophenyl)methyl]-4-[(oxazolo[4,5-d]pyridin-2-yl)amino]-1-piperidineethanol (412).

Example 34

- 5 A mixture of 2.7 parts of 2,3-dihydro-2-oxiranyl-1,4-benzodioxin, 3.5 parts of *N*-methyl-*N*-(4-piperidinyl)-2-benzoxazolamine, 45 parts of methylbenzene and 40 parts of methanol was stirred and refluxed overnight. The reaction mixture was evaporated and the residue was separated by column chromatography over
- 10 LiChroprep. RP 18 using a mixture of water, ammonium acetate and methanol (30:0.5:70 by volume) as eluent. The first fraction (A-isomer) was collected and the eluent was evaporated. The residue was taken up in water and the solution was treated with sodium hydroxide. The product was extracted with trichloromethane. The
- 15 extract was washed with water, dried, filtered and evaporated. The residue was crystallized from 2-propanol, yielding 0.3 parts (5%) of (A)-4-[(2-benzoxazolyl)methylamino]- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-1-piperidineethanol; mp. 160.4°C (413).
- The second fraction (B-isomer) was collected and the eluent was
- 20 evaporated. The residue was taken up in water and the solution was treated with sodium hydroxide. The product was extracted with trichloromethane. The extract was washed with water, dried, filtered and evaporated. The residue was crystallized from a mixture of 2,2'-oxybispropane and methanol, yielding 1.3 parts (21%) of (B)-4-[(2-benzoxazolyl)methylamino]- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-
- 25 1-piperidineethanol; mp. 120.9°C (414).

In a similar manner there were also prepared:

- (A)-4-[(2-benzothiazolyl)methylamino]- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-1-piperidineethanol; mp. 142.0°C (415).
- 30 (B)-4-[(2-benzothiazolyl)methylamino]- α -(2,3-dihydro-1,4-benzodioxin-2-yl)-1-piperidineethanol; mp. 134.2°C (416).

Example 35

- A mixture of 3.6 parts of [(3-chlorophenoxy)methyl]oxirane, 3.3 parts of *N*-methyl-*N*-(3-pyrrolidinyl)-2-benzoxazolamine and 90 parts
- 35 of methylbenzene was stirred and refluxed overnight. The reaction

mixture was evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (99:1 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted
 5 into the 4-methylbenzenesulfonate salt in 2-propanol. The salt was filtered off and dried, yielding 5 parts (58%) of 3-[(2-benzoxazoly1)methylamino]- α -[(3-chlorophenoxy)methyl]-1-pyrrolidine-ethanol 4-methylbenzenesulfonate(1:1); mp. 96.8°C (417).

In a similar manner there were also prepared:

- 10 4-[(2-benzothiazoly1)methylamino]- α -[(3-fluorophenoxy)methyl]-1-piperidineethanol dihydrochloride; mp. 205.2°C (418);
 4-[(2-benzothiazoly1)(phenylmethyl)amino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol (E)-2-butenedioate(2:1); mp. 163.5°C (419);
- 15 4-[methyl(6-methyl-2-benzothiazoly1)amino]- α -(phenoxymethyl)-1-piperidineethanol; mp. 140.6°C (420);
 α -[(4-fluorophenoxy)methyl]-4-[methyl(6-methyl-2-benzothiazoly1)amino]-1-piperidineethanol; mp. 105.5°C (421);
 α -[(3-fluorophenoxy)methyl]-4-[methyl(6-methyl-2-benzothiazoly1)amino]-1-piperidineethanol; mp. 109.6°C (422);
- 20 4-[(2-benzothiazoly1)(phenylmethyl)amino]- α -[(4-chlorophenoxy)methyl]-1-piperidineethanol (E)-2-butenedioate(2:1); mp. 172.0°C (423);
 4-[(2-benzothiazoly1)methylamino]- α -[(4-fluorophenoxy)methyl]- α -methyl-1-piperidineethanol 4-methylbenzenesulfonate(1:2); mp.
 25 207.3°C (424);
- cis-4-[(2-benzothiazoly1)methylamino]-3-hydroxy- α -(phenoxymethyl)-1-piperidineethanol; mp. 152.2°C (425);
cis-4-[(2-benzothiazoly1)amino]- α -[(4-fluorophenoxy)methyl]-3-hydroxy-1-piperidineethanol (Z)-2-butenedioate(1:2); mp. 156.7°C (426);
- 30 and
 4-[(2-benzothiazoly1)methylamino]- α -[[2,3-dihydro-1,4-benzodioxin-2-yl)-methoxy)methyl]-1-piperidineethanol (E)-2-butenedioate(1:1); mp. 146.5°C (427).

Example 36

- 35 A mixture of 2.77 parts of [(3-chlorophenoxy)methyl]oxirane, 4.92

parts of *N*-(3-methyl-2(3H)-benzothiazolylidene)-4-piperidinamine monohydrobromide, 2.12 parts of sodium carbonate, 60 parts of methanol and 68 parts of methylbenzene was stirred and refluxed overnight. The reaction mixture was cooled and the solvent was
5 evaporated. The residue was purified by filtration over silica gel using a mixture of trichloromethane and methanol (99:1 by volume), as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (Z)-2-butenedioate salt in 2-propanol and methanol. The salt was filtered off and
10 dried, yielding 5.5 parts (55%) of α -[(3-chlorophenoxy)methyl]-4-[(3-methyl-2(3H)-benzothiazolylden)amino]-1-piperidineethanol (Z)-2-butenedioate(1:2); mp. 157.6°C (428).

In a similar manner there were also prepared:

α -[(4-fluorophenoxy)methyl]-4-[(3-methyl-2(3H)-benzothiazolyl-
15 iden)amino]-1-piperidineethanol (Z)-2-butenedioate (1:2); mp. 140.2°C (429); and
3-[(2-benzothiazolyl)methylamino]- α -(phoxymethyl)-1-pyrrolidine-ethanol 4-methylbenzenesulfonate(1:2); mp. 196.3°C (430).

Example 37

20 A mixture of 2.5 parts of [(4-fluorophenoxy)methyl]oxirane, 4.3 parts of *N*-butyl-*N*-(4-piperidinyl)-2-benzothiazolamine, 45 parts of benzene and 40 parts of methanol was stirred and refluxed overnight. The reaction mixture was evaporated. The residue was purified by
25 column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (E)-2-butenedioate salt in ethanol. The salt was filtered off and dried, yielding 5.6 parts (72%) of 4-[(2-benzothiazolyl)butylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidine-
30 ethanol (E)-2-butenedioate(2:1); mp. 196.8°C (431).

In a similar manner there were also prepared:

4-[(2-benzothiazolyl)butylamino]- α -[(4-chlorophenoxy)methyl]-1-piperidineethanol (E)-2-butenedioate(2:1); mp. 192.2°C (432).

Example 38

A mixture of 1.8 parts of (B)-3,4-dihydro-2-oxiranyl-2H-1-benzopyran, 2.3 parts of N-methyl-N-(4-piperidinyl)-2-benzoxazoliné and 40 parts of ethanol was stirred and refluxed overnight. The reaction mixture was evaporated. The oily residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (90:10 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The oily residue was converted into the ethanedioate salt in 2-propanol. The salt was filtered off (slowly), washed with 2-propanol and 2,2'-oxybispropane and dried, yielding 3.82 parts (77.6%) of (B)-4-[(2-benzoxazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidine-ethanol ethanedioate (1:1); mp. 192.5°C (433).

Following the same procedure and using equivalent amounts of the appropriate starting materials, there were also prepared:

- (A)-4-[(2-benzothiazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol ethanedioate(1:1); mp. 195.0°C (434);
 (B)-4-[(2-benzothiazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol; mp. 133.4°C (435);
 (A)-4-[(2-benzoxazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol ethanedioate (1:1); mp. 206.0°C (436);
 A^+ -4-[(2-benzothiazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol ethanedioate(2:5); mp. 159.0°C (437);
 A^- -4-[(2-benzothiazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol; mp. 186.3°C; $[\alpha]_D^{25} = -4.61^\circ$ (c = 1% in methanol) (438); and
 A^+ -4-[(2-benzothiazolyl)methylamino]- α -(3,4-dihydro-2H-1-benzopyran-2-yl)-1-piperidineethanol; mp. 138.5°C; $[\alpha]^{25} = 47.60^\circ$ (c = 1% in methanol) (439).

30 Example 39

A mixture of 3.7 parts of [(phenylthio)methyl]oxirane, 5.0 parts of N-methyl-N-(4-piperidinyl)-2-benzothiazolamine and 120 parts of 2-propanol was stirred overnight at reflux temperature. The reaction mixture was evaporated. The residue was converted into the (Z)-2-butenedioate salt in 2-propanol and 2,2'-oxybispropane (5:1 by

volume). The salt was filtered off and dried, yielding 8.3 parts (78.3%) of 4-[(2-benzothiazolyl)methylamino]- α -[(phenylthio)methyl]-1-piperidineethanol (Z)-2-butenedioate(1:1); mp. 160.2°C (440).

5 In a similar manner there were also prepared:

4-[(2-benzothiazolyl)methylamino]- α -[(2-bromo-4-fluorophenoxy)methyl]-1-piperidineethanol; mp. 124.6°C (441);

4-[(2-benzothiazolyl)methylamino]- α -[[4-(4-fluorophenyl)thio]methyl]-1-piperidineethanol (Z)-2-butenedioate(1:1); mp. 140.6°C (442);

10 4-[(2-benzothiazolyl)methylamino]- α -[[4-(4-chlorophenyl)thio]methyl]-1-piperidineethanol (Z)-2-butenedioate(1:1); mp. 174.1°C (443);

4-[(2-benzoxazolyl)methylamino]- α -[[4-(4-fluorophenyl)thio]methyl]-1-piperidineethanol (Z)-2-butenedioate(1:1); mp. 141.7°C (444);

15 4-[(2-benzoxazolyl)methylamino]- α -[[4-(4-chlorophenyl)thio]methyl]-1-piperidineethanol (Z)-2-butenedioate(1:1); mp. 168.8°C (445);

4-[(2-benzoxazolyl)methylamino]- α -[(phenylthio)methyl]-1-piperidineethanol (E)-2-butenedioate(2:1); mp. 180.2°C (446); and α -[(4-fluorophenoxy)methyl]-4-[(thiazolo[4.5-c]pyridin-2-yl)amino]-1-piperidineethanol ethanedioate(1:2); mp. 195.9°C (447).

20 Example 40

A mixture of 1.65 parts of 1-(4-fluorobenzoyl)aziridine (solution in benzene 1.2M), 2.1 parts of N-(4-piperidinyl)-2-benzoxazolamine and 90 parts of methylbenzene was stirred and refluxed overnight. The reaction mixture was evaporated and the residue was crystallized
25 from 2-propanol. The product was filtered off and recrystallized from 2-propanone, yielding 1.3 parts (34%) of N-[2-[4-[(2-benzoxazolyl)amino]-1-piperidinyl]ethyl]-4-fluorobenzamide; mp. 173.3°C (448).

In a similar manner there were also prepared:

30 N-[2-[4-[2-(benzoxazolyl)methylamino]-1-piperidinyl]ethyl]-4-fluorobenzamide; mp. 162.9°C (449).

Example 41

A mixture of 3.2 parts of 4-(1-ethenyl)pyridine, 2.1 parts of N-(4-piperidinyl)-2-benzoxazolamine and 80 parts of 1-butanol was stirred and refluxed for 48 hours. The reaction mixture was
35 evaporated. The residue was purified by column chromatography over

silica gel using a mixture of trichloromethane and methanol (97:3 by volume) saturated with ammonia, as eluent. The pure fractions were collected and the eluent was evaporated. The residue was crystallized from 2,2'-oxybispropane, yielding 2 parts (62%) of N-[1-[2-(4-pyridinyl)ethyl]-4-piperidinyl]-2-benzoxazamine; mp. 146.9°C (450).

In a similar manner there were also prepared:

N-[1-[2-(2-pyridinyl)ethyl]-4-piperidinyl]-2-benzoxazamine; mp. 130°C (451); and

N-methyl-N-[1-[2-(2-pyridinyl)ethyl]-4-piperidinyl]-2-benzoxazamine; mp. 102.3°C (452).

Example 42

A mixture of 4.2 parts of 4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol, 4.1 parts of acetic acid anhydride, 0.1 parts of 4-methylbenzenesulfonic acid and 135 parts of methylbenzene was stirred and refluxed overnight. After cooling to room temperature, the reaction mixture was washed with a sodium hydroxide solution 5% and with water, dried, filtered and evaporated. The residue was crystallized from 2,2'-oxybispropane. The product was filtered off and dried, yielding 3.5 parts (77%) of 4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol acetate(ester); mp. 100.0°C (453).

Example 43

To a stirred and hot (60°C) solution of 4.2 parts of 4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol in 90 parts of N,N-dimethylformamide were added dropwise 2.3 parts of decanoyl chloride. Upon completion, stirring was continued for 24 hours at 60°C. After cooling to room temperature, the reaction mixture was poured onto water and the whole was treated with sodium hydroxide. The aqueous phase was decanted and the residual product was washed three times with water and dissolved in trichloromethane. The organic layer was dried, filtered and evaporated. The residue was purified by column chromatography over silica gel using a mixture of trichloromethane and methanol (97:3 by volume) as eluent. The pure fractions were collected and the eluent was evaporated. The residue was converted into the (Z)-2-butene-

dioate salt in a mixture of 2-propanone and 2,2'-oxybispropane. The salt was filtered off and dried, yielding 1.1 parts (16%) of [2-[4-[(2-benzothiazolyl)methylamino]-1-piperidinyl]-1-[(4-fluorophenoxy)methyl]ethyl]decanoate (Z)-2-butenedioate(1:1); mp. 134.2°C (454).

5 Example 44

3.4 Parts of (cis+trans)-N-[1-[4-(4-fluorophenoxy)cyclohexyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine were separated by column chromatography (HPLC) over silica gel using a mixture of trichloromethane and methanol (98:2 by volume) as eluent.

10 -The first fraction (A-isomer) was collected and the eluent was evaporated. The residue was crystallized from 2-propanol, yielding 0.34 parts of (A)-N-[1-[4-(4-fluorophenoxy)cyclohexyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 155.5°C (455).

-The second fraction (B-isomer) was collected and the eluent was
15 evaporated. The residue was crystallized from 2-propanol, yielding 0.34 parts of (B)-N-[1-[4-(4-fluorophenoxy)cyclohexyl]-4-piperidinyl]-N-methyl-2-benzothiazolamine; mp. 116.5°C (456).

Example 45

To a stirred solution of 5 parts of 4-[(2-benzothiazolyl)methyl-
20 amino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol in 80 parts of ethanol were added 1.8 parts of (+)-[R-(R*,R*)]-2,3-dihydroxybutanedioic acid while boiling. The product was allowed to crystallize. It was filtered off and dried, yielding 5.5 parts (94%)
25 of (+)-4-[(2-benzothiazolyl)methylamino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol [R-(R*,R*)]-2,3-dihydroxybutanedioate (2:1); mp. 189.1°C (457).

In a similar manner there were also prepared:

4-[(2-benzothiazolyl)methylamino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol 2-hydroxy-1,2,3-propanetricarboxylate(1:1);
30 mp. 146.7°C; (458) and

4-[(2-benzothiazolyl)methylamino]-α-[(4-fluorophenoxy)methyl]-1-piperidineethanol (Z)-2-butenedioate (1:1); mp. 127.6°C (459).

C. Pharmacological Examples

The useful anti-anoxic properties of the compounds of formula (I) can be demonstrated in the following test procedures.

5 Example 46:KCN Test in Rats.

Histotoxic anoxia is produced by a rapid intravenous injection of potassium cyanide (KCN) in rats.

10 KCN in control rats induces abdominal contractions and clonic seizures and is lethal within 2 minutes. The protection from KCN induced lethality appears to be a simple way of evaluating quantitatively the anti-anoxic properties of the test compounds. Male Wistar rats weighing \pm 200 g were treated subcutaneously with a test compound. One hour after treatment there was injected intravenously a KCN solution at a dose of 5
15 mg/kg body weight. Survival after 2 minutes was evaluated and ED_{50} -values were determined.

20 The ED_{50} -values of the compounds of formula (I) are listed in table I. Said ED_{50} -values are those values in mg/kg body weight capable of protecting 50% of the tested animals against KCN-induced lethality.

25 The compounds listed in table I are not given for the purpose of limiting the invention thereto but only to exemplify the useful pharmacological activities of all the compounds within the scope of formula (I).

Compound number	ED_{50} -values in mg/kg body weight
1	5
9	2.5
10	5
11	1.25
15	0.63

	18	5
	24	5
	31	5
5	32	5
	42	2.5
	48	1.25
	50	2.5
	75	10
10	103	5
	107	2.5
	108	5
	118	5
	120	2.5
15	121	2.5
	123	5
	133	5
	146	5
	148	5
20	151	5
	184	5
	191	2.5
	195	1.25
	309	5
25	320	1.8
	322	5
	323	1.25
	324	2.5
	357	0.63
30	405	10
	410	5
	411	5
	413	5

5

10

415	10
438	10
440	10
442	2.5
443	10
446	10
452	5

D) Composition Examples.

The following formulations exemplify typical pharmaceutical compositions in dosage unit form suitable for systemic administration to animal and human subjects in accordance with the instant invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I) or pharmaceutically acceptable acid addition salt thereof.

Example 47 : ORAL DROPS

500 Grams of the A.I. was dissolved in 0.5 liters of 2-hydroxypropanoic acid and 1.5 liters of the polyethylene glycol at 60-80°C. After cooling to 30-40°C there were added 35 liters of polyethylene glycol and the mixture was stirred well. Then there was added a solution of 1750 grams of sodium saccharin in 2.5 liters of purified water and while stirring there were added 2.5 liters of cocoa flavor and polyethylene glycol q.s. to a volume of 50 liters, providing an oral drop solution comprising 10 milligrams of the A.I. per milliliter. The resulting solution was filled into suitable containers.

Example 48 : ORAL SOLUTION

9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 liters of boiling purified water. In 3 liters of this solution were dissolved first 10 grams of

2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 liters 1,2,3-propanetriol and 3 liters of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved
5 in 0.5 liters of water and 2 milliliters of raspberry and 2 milliliters of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 liters providing an oral solution comprising 20 milligrams of the active ingredient per teaspoonful (5 milliliters). The resulting solution was filled in
10 suitable containers.

Example 49 : CAPSULES

20 Grams of the A.I., 6 grams sodium lauryl sulfate, 56 grams starch, 56 grams lactose, 0.8 grams colloidal silicon dioxide, and 1.2 grams magnesium stearate were vigorously stirred together. The resulting
15 mixture was subsequently filled into 1000 suitable hardened gelating capsules, comprising each 20 milligrams of the active ingredient.

Example 50 : FILM-COATED TABLETS

Preparation of tablet core

A mixture of 100 grams of the A.I., 570 grams lactose and 200 grams
20 starch was mixed well and thereafter humidified with a solution of 5 grams sodium dodecyl sulfate and 10 grams polyvinylpyrrolidone (Kollidon-K 90®) in about 200 milliliters of water. The wet powder mixture was sieved, dried and sieved again. Then there was added 100 grams microcrystalline cellulose (Avicel®) and 15 grams hydrogenated
25 vegetable oil (Sterotex ®). The whole was mixed well and compressed into tablets, giving 10,000 tablets, each containing 10 milligrams of the active ingredient.

Coating

To a solution of 10 grams methyl cellulose (Methocel 60 HG®) in
30 75 milliliters of denaturated ethanol there was added a solution of 5 grams of ethyl cellulose (Ethocel 22 cps ®) in 150 milliliters of dichloromethane. Then there were added 75 milliliters of dichloromethane and 2.5 milliliters 1,2,3-propanetriol. 10 Grams of polyethylene glycol was molten and dissolved in 75 milliliters of dichloromethane. The
35 latter solution was added to the former and then there were added 2.5

grams of magnesium octadecanoate, 5 grams of polyvinylpyrrolidone and 30 milliliters of concentrated colour suspension (Opaspray K-1-2109●) and the whole was homogenated.

5 The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example 51 : INJECTABLE SOLUTION

1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 liters of boiling water for injection. After cooling to about 50°C there were added while stirring 4
10 grams lactic acid, 0.05 grams propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 liter volume, giving a solution of 4 milligrams A.I. per milliliters. The solution was sterilized by filtration (U.S.P. XVII p. 811) and filled in sterile containers.

15 Example 52 : SUPPOSITORIES


3 Grams A.I. was dissolved in a solution of 3 grams 2,3-dihydroxybutanedioic acid in 25 milliliters polyethylene glycol 400. 12 Grams surfactant (SPAN●) and triglycerides (Witepsol 555 ●) q.s. ad 300
20 grams were molten together. The latter mixture was mixed well with the former solution. The thus obtained mixture was poured into moulds at a temperature of 37-38°C to form 100 suppositories each containing 30 milligrams of the active ingredient.

25

30

35

1. A chemical compound having the formula



4 $-A^1=A^2-A^3=A^4-$ is a bivalent radical having the formula

6 $-N=CH-CH=CH-$ (b):

8 $-\text{CH}=\text{CH}-\text{N}=\text{CH}-$ (d); or

9 $-\text{CH}=\text{CH}-\text{CH}=\text{N}-$ (e):

10 wherein one or two hydrogen atoms in the radical $-A^1=A^2-A^3=A^4-$

11 may, each independently from each other, be replaced by halo, C_{1-6}

12 alkyl, hydroxy, C₁₋₆ alkyloxy or trifluoromethyl;

13 Z is 0 or S;

14 n is 0 or the integer 1:

15 R is a member selected from the group consisting of hydrogen,

16 C₁₋₆ alkyl, hydroxy and C₁₋₆ alkyloxy or trifluoromethyl;

17 R^1 or R^2 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

18 the dotted line between the nitrogen atom bearing R¹, the

19 interjacent carbon atom and the nitrogen atom bearing R²

20 indicating that a double bond exists between the nitrogen bearing

21 R¹ and said interjacent carbon atom, in which case R¹ is absent,

22 or a double bond exists between the interjacent carbon and the

23 nitrogen bearing R^2 , in which case R^2 is absent; and wherein

24 said interjacent carbon atom is the carbon atom positioned between

25 the nitrogen bearing R¹ and the nitrogen bearing R²;



26 R^3 or R^4 are both hydrogens or R^3 and R^4 combined may

27 form a bivalent radical of formula $-\text{CH}_2-\text{CH}_2-$;

28 L is a member selected from the group consisting of C_{1-12} alkyl,

29 substituted C_{1-6} alkyl, aryl C_{2-6} alkenyl, C_{3-6} cycloalkyl,

30 substituted C_{3-6} cycloalkyl, a radical of formula



33 R^5 being hydrogen or C_{1-12} alkylcarbonyl;

34 R^6 being hydrogen or C_{1-6} alkyl;

35 R^7 being aryl, a radical R^8-O or a radical R^8-S ;

36 B being a bivalent radical of formula $-\text{CH}_2-$ or $-\text{O}-$; and

37 said R^8 being hydrogen; aryl; 2,3-dihydro-1H-indenyl;

38 benzodioxolyl; (2,3-dihydro-1,4-benzodioxin-2-yl)methyl;

39 (2H-1-benzopyran-2-yl)methyl; or phenyl substituted with C_{2-6}

40 alkenyloxy, C_{1-6} alkylcarbonylamino or C_{1-6} alkylphenylcarbonyl; and

41 wherein said substituted C_{3-6} cycloalkyl is C_{3-6} cycloalkyl

42 being substituted with up to two substituents each independently

43 selected from the group consisting of aryl, aryloxy and cyano;

44 said substituted C_{1-6} alkyl is C_{1-6} alkyl being substituted

45 with a member selected from the group consisting of C_{3-6} cycloalkyl,

46 pyridinyl, pyridinyloxy, aryl, benzimidazolyl, indolyl, isoxazolyl

47 being optionally substituted with phenyl, 3-oxo-1,2,4-triazolo[4,5-a]-

48 pyridin-2(3H)-yl, a radical of formula $\text{ArY}-$, a radical of formula

49 $\text{R}^9\text{O}-$ and a radical of formula



51 said Ar being aryl;

52 Y being O, S, NH, $-\text{CONH}-$, $-\text{CHCN}$, $-\text{CHOR}^{10}$ or $-\text{SO}_2$;

53 R^9 being hydrogen, aryl C_{1-6} alkyl, C_{3-6} cycloalkyl, phenyl

54 substituted with phenyl, or phenyl substituted with C_{1-6} alkyloxy-

carbonyl;

R^{10} being hydrogen or C_{1-6} alkyl; and

wherein aryl is phenyl optionally substituted with up to three substituents each independently selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkyloxy, halo, trifluoromethyl, cyano, C_{1-6} alkylcarbonyl, nitro, amino and aminocarbonyl.

2. A chemical compound according to claim 1, wherein L is aryl C_{2-6} alkenyl, C_{1-6} alkyl being substituted with aryloxy, a radical of formula (a) or a radical of formula (b).

3. A chemical compound according to claim 2, wherein $-A^1=A^2-A^3=A^4-$ is $-CH=CH-CH=CH-$, n is the integer 1, R^2 is absent and a double bond exists between the nitrogen bearing R^2 and the interjacent carbon atom, and R^3 and R^4 are both hydrogen atoms.

4. A chemical compound according to claim 3 wherein in the radical of formula (a), R^5 and R^6 are both hydrogen and R^7 is a radical R^8-O- , wherein R^8 is aryl; or wherein in the radical of formula (b), B is O and R^5 and R^6 are both hydrogen radicals.

5. A chemical compound according to claim 4 wherein R^1 is C_{1-6} alkyl and aryl is halophenyl.

6. A chemical compound according to claim 1, wherein the compound is 4-[(2-benzothiazolyl)methylamino]- α -[(4-fluorophenoxy)methyl]-1-piperidineethanol.

7. A pharmaceutical composition comprising a suitable pharmaceutical carrier and as an active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6.

8. An anti-anoxic or anti-hypoxic composition comprising a suitable pharmaceutical carrier and as an active ingredient an effective anti-anoxic or anti-hypoxic amount of a compound as claimed in any one of claims 1 to 6.

9. A method of preparing a pharmaceutical composition, characterized in that a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6 is intimately mixed with suitable pharmaceutical carriers.


10. A compound as claimed in any one of claims 1 to 6 for use as a medicine.

BAD ORIGINAL


1 11. A compound as claimed in any one of claims 1 to 6 for use as an
2 anti-anoxic or anti-hypoxic medicine.

1 12. A process for preparing a chemical compound as claimed in
2 in claim 1, characterized by

3 a) N-alkylating a piperidine of formula



4 with a reagent of formula




5 wherein W represents a reactive leaving group, in a reaction-
6 inert solvent;

7 b) reductively N-alkylating a piperidine of formula (III) with a
8 carbonyl-compound of formula




9 whereby L'=C=O is a compound of formula L-H wherein a $-\text{CH}_2-$
10 radical is oxidized to a carbonyl radical, in a reaction-inert
11 solvent;

12 c) cyclizing an urea or thiourea derivative of formula




13 in a reaction-inert solvent in the presence of an oxidant;

14 d) reacting a piperidine of formula (III) with an oxirane of formula




in a reaction-inert solvent, thus preparing a compound of formula




16 and, if desired, acylating (I-a-1) with a reagent




17 said $\text{R}^{5-a}\text{-W}^1$ representing an alkanolic acid or a reactive
18 derivative thereof, thus preparing a compound of formula




19 e) reacting a piperidine of formula (III) with an oxirane of formula




20 in a reaction-inert solvent, thus preparing a compound of formula




21 and, if desired, acylating (I-a-3) with a reagent (VII), thus
22 preparing a compound of formula




23 f) reacting a benzothiazole or benzoxazole of formula




24 with a reagent of formula




25 in a reaction-inert solvent, wherein Q^1 is a reactive leaving
 26 group and Q^2 is a radical of formula $-NHR^1$, or alternatively,
 27 Q^1 is a radical of formula $-NHR^1$ and Q^2 is a reactive
 28 leaving group, thus preparing a compound of formula




29 g) cyclodesulfurizing a thiourea derivative of formula




30 which in situ may be prepared by condensing an isothiocyanate




31 with an aromatic aminoalcohol of formula




32 thus preparing a compound of formula




33 h) cyclizing an urea or thiourea derivative of formula




34 which in situ may be formed by reacting an isocyanate or
35 isothiocyanate of formula



36 with an aromatic amine of formula



37 wherein W² represents a leaving group, thus preparing a compound
38 of formula




39 1) reacting a piperidine of formula (III) with an alkene derivative

40 of formula



41 in a reaction-inert solvent, thus preparing a compound of formula




42 wherein R^{11} represents C_{1-6} alkyl, C_{3-6} cycloalkyl,
43 pyridinyl, pyridinyloxy, aryl, benzimidazolyl, indolyl,
44 isoxazolyl being substituted with phenyl, 3-oxo-1,2,4-triazolo-
45 [4,5-a]pyridin-2(3H)-yl, a radical of formula ArY , a radical of
46 formula R^9O and a radical of formula (g); and r is an integer
47 of from 2 to 6 included.


48 j) reacting a piperidine of formula (III) with an aziridine of
49 formula



50 in a reaction-inert solvent thus preparing a compound of formula



51 wherein D represents a radical of formula



52 and, if desired, converting the compounds into each other following
53 art-known functional groups-transformation reactions; and if further
54 desired, converting the compounds of formula (I) into a therapeutically
55 active non-toxic acid-addition salt form by treatment with an
56 appropriate acid or, conversely, converting the acid-addition salt
57 into the free base form with alkali; and/or preparing stereochemically
58 isomeric forms thereof.



European Patent
Office

EUROPEAN SEARCH REPORT

0184257

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 85201937.1
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	DE - A1 - 2 545 645 (BAYER) * Formula I * --	1	C 07 D 413/12 C 07 D 417/12 C 07 D 413/14
A	CHEMICAL ABSTRACTS, vol. 97, no. 5, August 2, 1982, Columbus, Ohio, USA MARCHALIN, MIROSLAV; POVAZANEC, FRANTISEK; MARTVON, AUGUSTIN "A study of the reaction of 2-pyridyl isothiocyanate with azoimide." page 570, column 2, abstract no. 38 892h & Collect. Czech. Chem. Commun. 1982, 47(3), 877-83 --	1	C 07 D 417/14 C 07 D 498/04 C 07 D 513/04 A 61 K 31/445 A 61 K 31/42 A 61 K 31/425
A	CHEMICAL ABSTRACTS, vol. 78, no. 25, June 25, 1973, Columbus, Ohio, USA HISANO, TAKUZO; YABUTA, YUKO "Synthesis of organosulfur compounds. VIII. Cyclization products from the modified Willgerodt-Kindler reaction." page 408, column 1, abstract no. 159 505a & Chem. Pharm. Bull. 1973, 21(3), 511-17 --	1	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 07 D 413/00 C 07 D 417/00 C 07 D 498/00 C 07 D 513/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 17-01-1986	Examiner HAMMER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



European Patent
Office

EUROPEAN SEARCH REPORT

0184257

Application number

EP 85201937.1

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	<p>CHEMICAL ABSTRACTS, vol. 90, no. 21, May 21, 1979, Columbus, Ohio, USA</p> <p>SIMOV, D.; DAVIDKOV, K. "Valence vibrations of the azomethine group of 2-aminobenzoxazole derivatives." page 518, column 2, abstract no. 167 524u</p> <p>& Khim. Geterotsikl. Soedin. 1979, (2), 188-90</p> <p>--</p>	1	
A	<p>CHEMICAL ABSTRACTS, vol. 98, no. 21, May 23, 1983, Columbus, Ohio, USA</p> <p>MERCHAN, F.L.; GARIN, J.; MELENDEZ, E.; TEJERO, T. "2-(2-Benzimidazolylamino)benzothiazoles and 2-(2-imidazolidinylidenamino)benzothiazoles." page 651, column 1, abstract no. 179 281r</p> <p>& Synthesis 1982, (12), 1066-7</p> <p>--</p>	1	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 17-01-1986	Examiner HAMMER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

EUROPEAN SEARCH REPORT

0184257

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 85201937.1
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	CHEMICAL ABSTRACTS, vol. 98, no. 13, March 28, 1983, Columbus, Ohio, USA AHUJA, PADMA; SINGH, JUJHAR; NIGAM, M.B.; SARDANA, VINOD; KAR, K.; ANAND, NITYA "Lactam acetals. Part VII. Synthesis and biological activities of some heterocyclic amidines and diamidines." page 584, column 1, abstract no. 107 106q & Indian J. Chem., Sect. 1982, 21B(9), 849-52 --	1	
A	CHEMICAL ABSTRACTS, vol. 99, no. 7, August 15, 1983, Columbus, Ohio, USA SINGH S.P.; PRAKASH, INDRA; TOMER, R.K.; PRAKASH, O.M.; SAWHNEY, S.N. "Synthesis and mass spectra of some substituted 2-(2'-benzazolylamino)pyrimidines." page 553, column 2, abstract no. 53 691b & Indian J. Chem., Sect. B 1983, 22B(1), 37-42 --	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
A	DE - A - 1 804 450 (EGYESÜLT GYORGYSZER-...) * Claims 1,2,7,10 * --	1,7,9, 10,12,	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 17-01-1986	Examiner HAMMER
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



European Patent
Office

EUROPEAN SEARCH REPORT

0184257

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 85201937.1
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	<p>US - A - 2 623 878 (ISLER..)</p> <p>* Example 1; column 1, lines 36,37,52-54 *</p> <p>-----</p>	1,7,9,10	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
VIENNA		17-01-1986	HAMMER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.